

Long-Term Monitoring of Coral Reefs of the Main Hawaiian Islands

Final Report

2009 NOAA Coral Reef Conservation Program

Hawai'i Island Monitoring Report

NA09NOS4260100

10/01/2009 – 12/31/2012

William Walsh, Stephen Cotton, Camille Barnett, Linda
Preskitt, Courtney Couch



***Department of Land & Natural Resources
December 2012***

Contents

SUMMARY OF FINDINGS	7
CONTRIBUTORS	10
HAWAI'I ISLAND SURVEYS	12
Benthic Monitoring Methods	11
Results	11
Octocoral Distribution	14
Coral Disease	16
Field Surveys	16
Colony Assessment	17
Coral disease by size class	17
Coral community structure	18
Coral diseases in Hawai'i	19
Disease distribution and prevalence	20
Spatial patterns	22
Temporal comparisons	24
Temperature data	25
Fish Surveys	27
Shallow water resource fish surveys	29
Adult yellow tang surveys	30
Depth stratified random surveys	30
Retrospective surveys	31
Fish Surveys Results-West Hawai'i	32
Introduced Species – Ta'ape	35
Roi	36
Pufferfish Die-Off	39
Roi Impacts	40
Aquarium Species	42
Fish Replenishment Areas (FRAs)	46
Aquarium Species of Special Concern	54
Aquarium Species Open vs. FRA Trend Analysis	55
Aquarium Species Population and Catch Analysis	63
Endemic Species on the WHFC White List	66

Aquarium reef fish catch vs. non-aquarium catch	67
Gill Net Management.....	72
Invertebrates – Crown-of-Thorns.....	74
Crown-of-Thorns Aggregation Monitored on Kona Coast	74
Urchins.....	79
East Hawai'i Fish Survey Results	80
References.....	85

Figures

Figure 1. Comparison between survey years of percent coral cover across West Hawai'i monitoring sites.....	13
Figure 2. Same location at Puako showing significant loss of <i>Porites lobata</i>	14
Figure 3. <i>Sarcothelia edmonsoni</i> and another unidentified octocoral found on West Hawai'i reefs s	15
Figure 4. Comparison between survey years of percent cover of octocoral across West Hawai'i monitoring sites	16
Figure 5. Size structure of coral colonies recorded in West Hawai'i during surveys conducted in 2010.....	18
Figure 6. Examples of coral diseases observed in West Hawai'i during 2010 baseline surveys	20
Figure 7. Relative abundance of coral diseases recorded for DAR monitoring sites in West Hawai'i during surveys conducted in 2010	21
Figure 8. Prevalence of Poritid diseases at each West Hawai'i site in 2010	22
Figure 9. Disease prevalence in relation to site distance from harbors/boat ramps in West Hawai'i.....	23
Figure 10. Comparison between survey years (2007 vs. 2010) of diseased colony densities for three types of Poritid conditions at 10 sites in West Hawai'i	24
Figure 11. Temperature (°F) records for three West Hawai'i FRAs.....	26
Figure 12. West Hawai'i monitoring sites.....	27
Figure 13. Map showing locations of West Hawai'i shallow water resource fish surveys and laynet prohibited areas.....	29
Figure 14. Map showing the locations of Keauhou stratified random fish population survey sites.....	31
Figure 15. Overall temporal trend in mean fish density of three major fish utilization groups at West Hawai'i sites	32
Figure 16. Temporal trend in mean fish density for four families on West Hawai'i sites	34

Figure 17. Temporal trend in mean fish density for various fish species of interest at West Hawai'i sites.....	35
Figure 18. Overall change in herbivore biomass in FRA, MPA and open areas 2003-2012.....	35
Figure 19. Ta'ape abundance on West Hawai'i Transects and 10 minute free swim surveys	36
Figure 20. <i>C. argus</i> density trends in West Hawai'i.....	38
Figure 21. Deep-Sea swallower, <i>Kali indica</i> with inflated swim bladder	38
Figure 22. Number of dead puffers examined as USGS Honolulu Field Station	39
Figure 23. Pufferfish abundance trends in West Hawai'i.....	40
Figure 24. Temporal trends of the numbers of kole, yellow tang and roi in FRAs	41
Figure 25. Relationship between Roi and various West Hawai'i fish population parameters.....	42
Figure 26. Number of aquarium animals collected and number of commercial aquarium permits in West Hawai'i for fiscal years 1976-2012.....	43
Figure 27. Opae ula catch in East and West Hawai'i	44
Figure 28. Comparison between Hawai'i Island aquarium collector report data and dealer purchases of aquarium animals from the collectors (FY 2010).....	45
Figure 29. Effectiveness of individual FRAs to replenish yellow tang, 1999-2009.....	48
Figure 30. Overall changes in yellow tang abundance (Mean \pm SE) in FRAs, MPAs and Open areas, 1999-2012	49
Figure 31. Yellow tang abundance in West Hawai'i shallow water (10'-20') habitat	49
Figure 32. Overall changes in goldring surgeonfish (kole) abundance in FRAs, MPAs and Open areas, 1999-2012	50
Figure 33. Overall changes in Achilles tang abundance in FRAs, MPAs and Open areas, 1999-2012.....	51
Figure 34. Overall changes in orangespine unicornfish abundance in FRAs, MPAs and Open areas, 1999-2012	52
Figure 35. Overall changes in chevron tang abundance in FRAs, MPAs and Open areas, 1999-2012.....	53
Figure 36. Flame angelfish on transects in FRAs, MPAs and Open areas, 1999-2012.....	54
Figure 37. Flame angelfish on Free Swim surveys in FRAs, MPAs and Open areas, 1999-2012.....	54
Figure 38. White list species showing fairly consistent lower densities in areas open to aquarium collecting	58
Figure 39. White list species exhibiting inconsistent differences in density between areas open to aquarium collecting and FRAs.....	59
Figure 40. White list species exhibiting higher population densities in areas open to collecting relative to FRAs.....	61

Figure 41. West Hawai'i Yellow tang catch since FRA establishment and price per fish.....	65
Figure 42. Comparison of the number of reef fishes caught by recreational and commercial fishers in the Main Hawaiian Islands	68
Figure 43. Comparison of the biomass of reef fishes caught by recreational and commercial fishers in the Main Hawaiian Islands	69
Figure 44. Comparison of the number of reef fishes caught by recreational, commercial and aquarium fishers in West Hawai'i	69
Figure 45. Comparison of the no. of reef fishes, excluding yellow tang, caught by recreational, commercial and aquarium fishers in West Hawai'i.....	70
Figure 46. Comparison of the biomass of reef fishes caught by recreational, commercial and aquarium fishers in West Hawai'i	71
Figure 47. Comparison of the biomass of reef fishes, excluding yellow tang, caught by recreational, commercial and aquarium fishers in West Hawai'i.....	71
Figure 48. Comparison of the numbers of reef fishes caught by recreational, commercial and aquarium fishers on Maui.....	72
Figure 49. Comparison of the biomass of reef fishes caught by recreational, commercial and aquarium fishers on Maui.....	72
Figure 50. Locations of laynet prohibited areas in West Hawai'i and shallow water resource fish survey sites.....	73
Figure 51. Biomass of 'Resource' (i.e. food) fish on shallow water transects.....	74
Figure 52. Overall Crown-of-Thorns abundance on West Hawai'i transects and 10 minute free swim surveys.....	75
Figure 53. <i>Acanthaster planci</i> (COTS) density and abundances recorded on surveys conducted near Ka'ūpūlehu	76
Figure 54. Number of coral colonies observed at WHAP Site 7 with COTS feeding 'scars' and approximate area of subsequent colony mortality	77
Figure 55. Crown-of-Thorns Starfish (COTS) aggregation monitored at Ka'ūpūlehu in September 2012	79
Figure 56. Abundance (<i>Mean ± SE</i>) of collector urchin <i>Tripneustes gratilla</i> , red pencil urchin <i>Heterocentrotus mammillatus</i> and banded urchin <i>Echinothrix calamaris</i> on West Hawai'i transects.....	80
Figure 57. Fish community parameters at Waiopae (MLCD & OPEN) and Richardson's Ocean Center	81
Figure 58. Annual mean abundance of fishes at Waiopae and Richardson's Ocean Center.....	82
Figure 59. Wai Opae Open/MLCD and ROC fish abundance by family (all years pooled)	84

Tables

Table 1. Percent coral cover at West Hawai'i sites in 2003, 2007, 2011	12
Table 2. Occurrence of diseases across ten monitoring stations in survey years 2007 and 2010 in West Hawai'i	20
Table 3. West Hawai'i monitoring sites with corresponding coordinates, status and survey type	28
Table 4. List of fishes collected or reported in West Hawai'i die-off	37
Table 5. Changes in West Hawai'i aquarium fishery since implementation of the FRAs	43
Table 6. Overall FRA effectiveness for the top ten most aquarium collected fishes	47
Table 7. Proposed 'White List' of species which can be taken by aquarium collectors within the West Hawai'i Regional Fisheries Management Area	55
Table 8. Population estimates and % of population taken by aquarium collectors of 'White List' species	64
Table 9. . Endemic species on 'white list'	66
Table 10. Comparison of the number and pounds of reef fishes caught by recreational and commercial fishers on Maui and in West Hawai'i	72
Table 11. Percent Similarity from pairwise East Hawai'i site comparisons	81
Table 12. Fish Density per 100 m ² by family at East Hawai'i sites	83

Appendices

Appendix A. Occurrences of eight coral diseases documented across 30 monitoring sites in West Hawai'i	88
Appendix B. West Hawai'i Benthic Cover 2011 Surveys; Broad Benthic Categories	89
Appendix C. West Hawai'i Coral Cover By Species 2011 Surveys	90
Appendix D. West Hawai'i Benthic Cover 2007 Surveys; Broad Benthic Categories	91
Appendix E. West Hawai'i Coral Cover By Species 2007 Surveys	92
Appendix F. West Hawai'i Benthic Cover 2003 Surveys; Broad Benthic Categories	93
Appendix G. West Hawai'i Coral Cover By Species 2003 Surveys	94
Appendix H. 2003, 2007 and 2011 Octocoral Percent Cover Comparison	95
Appendix I. Individuals per 100 m ² by species at East Hawai'i sites	96

SUMMARY OF FINDINGS

Hawai'i

Benthic

Coral and Habitat Surveys

- Total Coral cover declined significantly at 6 northern sites in West Hawai'i between 2003 and 2007. A strong winter storm in 2004 was likely responsible for the declines but a major sediment event in 2006 may also have affected sites at Kamilo Gulch and Waiaka'ilio Bay on the North Kohala coast.
- No invasive alien algal or coral species were detected at any site. Macroalgal cover was very low at all sites.
- The distribution of the octocoral *Sarcothelia edmonsoni* around developed areas near Kona and its virtual absence around undeveloped shoreline areas suggests possible anthropogenic (pollution) influence. Since other studies have cited octocoral as a pollution indicator and shoreline development in West Hawai'i is expected to continue to increase, further studies should be undertaken to determine the relationship between octocoral presence and land based pollution.

Coral Disease Surveys

- The following coral diseases were recorded at West Hawai'i monitoring sites in 2010: *Porites* growth anomaly, *Porites* tissue loss syndrome, *Porites* multifocal tissue loss, *Porites* trematodiasis, *Montipora* growth anomaly, *Pavona varians* hypermycosis, *Pocillopora* tissue loss.
- *Porites* spp. were the most susceptible to disease (mean prevalence of $3.76 \pm 3.58\%$), with the most widespread diseases including growth anomalies, trematodiasis, and tissue loss syndrome of *Porites* spp.
- Though thought to be a common condition, the possible senescence reaction of *Pocillopora meandrina* (i.e. progressive age-related colony death) was observed at only two sites likely attributed to the low number of Pocilloporids present at monitoring sites.
- Overall disease prevalence and prevalence of *Porites* growth anomalies were positively correlated with total estimated size and total number of submarine groundwater (SGD) "plumes".
- West Hawai'i sites show a significant negative relationship between disease prevalence and distance from harbors/boat ramps (overall disease prevalence: $r = -0.402$, $p = 0.028$), particularly for *Porites* growth anomalies ($r = -0.658$, $p = 0.000$) and *Porites* tissue loss syndrome ($r = -0.701$, $p = 0.000$).
- No significant changes in disease densities were found between survey years 2007 and 2010 for ten DAR monitoring sites ($p = 0.18$). However, instances of *Porites* growth anomalies and *Porites* tissue loss syndrome slightly increased at four sites located in close proximity to harbors/boat ramps.
- No statistically significant relationships were found between prevalence of coral diseases and abundances of corallivorous butterflyfishes and parrotfishes for West Hawaii's reefs.

Fish

- The abundance of both aquarium and food fishes increased significantly in West Hawai'i over the last 11 years. The overall number of fishes not substantially harvested for either food or for the aquarium trade, did not change significantly although individual species within this group may have.
- Examination of the temporal trends of some of the most common reef fish families indicates that acanthurids have been increasing over the past eleven years while labrids have decreased. Overall, chaetodontids and pomacentrids have been relatively stable although some species within the family have either increased or declined.

Introduced Species/Fish Die-Off

- Transect data reflects overall low abundance of ta'ape in the reef areas of the study sites and they are rarely found in the shallower water where resource fish surveys are conducted. Ta'ape are numerous in some locales usually along drop-offs and deeper reef areas but their distribution is highly patchy and they are not at all abundant in many reef areas in West Hawai'i. Ta'ape numbers also appear to have declined from earlier periods.
- There has been a marked decrease in roi abundance both on West Hawai'i transect (56% decrease) and free swim surveys (55% decrease). This decline may be related in part to an unusual fish die-off in West Hawai'i which first became apparent in May 2006.
- Early in 2010 a die-off of large puffers, with external symptoms quite similar to the previous mortalities, began to occur on Maui and Hawai'i Island. Over the ensuing months low numbers of dead and dying puffers were progressively reported up the island chain as far as Kaua'i (Oct. 2010).
- West Hawai'i monitoring data indicates a substantial decline has occurred in the abundance of the Hawaiian spotted toby (*Canthigaster jactator*) and the spotted puffer (*Arothron meleagris*) with a precipitous drop of the latter species in 2009/2010.
- As of November 2010 a total of 106 puffers have undergone both gross and microscopic examination. All assays for viruses (including electron microscopy) have so far come up negative and all attempts to incriminate any infectious agent as a cause have come to naught.
- An examination of roi and two of the most abundant species in roi's prime habitat the yellow tang (*Zebrasoma flavescens*) and kole (*Ctenochaetus strigosus*) fails to indicate direct negative impact on either species.
- Examination of the relationship between roi abundance and the abundance of various species and functional groups shows no significant negative relationships. In other words having more roi in an area does not result in having less total fish, small prey fish, other piscivores, yellow tang Young-of-Year (YOY), kole YOY or all YOY.
- The estimated roi population in West Hawai'i in the 30'-60' depth range (hard bottom only) is 58,839 individuals.

Aquarium Species

- Ten years after closure of the FRAs, the top 20 aquarium species showed a small overall increase in abundance relative to the period before the FRAs were operational. Most of the increase was attributed to the top two species Yellow Tang and Goldring Surgeonfish (kole) which comprise 91% of the West Hawai'i Aquarium catch.
- Seven of the top 10 most collected species (representing <6% of all collected fish) decreased in overall density. Three of these decreases were significant (Achilles Tang, Multiband Butterflyfish and Black Surgeonfish).
- The FRAs were 'effective' (increases in FRAs relative to long-term MPAs) for eight of the top 10 collected species with three being statistically significant. With only a single exception the FRAs were highly effective in increasing the abundance of Yellow Tang along the West Hawai'i coastline.
- A decrease of yellow tang in open areas to below baseline levels is largely attributable to an increase in the number of aquarium collectors and collected animals relative to the period when the FRAs were established. Kole are more abundant and much less collected than the yellow tangs and open areas have been relatively stable. Achilles tangs are in a declining trend.
- Concerns over continued expansion of the aquarium fishery and harvesting effects in the open areas has prompted DAR and the West Hawai'i Fisheries Council (WHFC) to develop a 'white list' of 40 species which can be taken by aquarium fishers. All other species are off limits.
- Based on an analysis of the differences in density between open and protected areas there was clear evidence of an aquarium collecting impact for only 5 species of the 34 white list species which were analyzed. Four of the 5 are among the 10 most heavily collected species. For the others, it appears that inclusion on the white list poses little or no threat to their populations.
- Based on a comparison of catch and estimated population abundance in the 30'-60' depth range aquarium collecting is having the largest impacts on Achilles and yellow tang. Achilles tang has had low levels of recruitment over the past decade and substantial numbers of larger fish (i.e. 'breeders') are taken for human consumption. Yellow tang has generally recruited reliably but aquarium take has risen dramatically to currently unsustainable levels.
- For most of the species on the white list collecting impact, in terms of the % of the population being removed annually, is relatively low with 10 species having single digit % catch and 19 species having % catch values <1%.
- Eight no lay gill netting areas were established in West Hawai'i in 2005, comprising 25% of the coastline (including already protected areas). Nearshore monitoring results did not find major differences in food fish abundance in/out of the no netting areas. The lack of a marked effect of protection may be due to several factors including the relatively low number of lay gill nets that are presently being used (i.e. registered) in West Hawai'i.

Invertebrates

- Crown-of-Thorns Starfish (COTS) (*Acanthaster planci*) have a low absolute abundance on West Hawai'i reefs and there has been an overall decreasing trend in abundance over the last four years.

- An aggregation of COTS was monitored near Ka'ūpūlehu, West Hawai'i at WHAP Site 7 in September 2012. COTS were noticeably distributed in clusters.
- Surveys conducted at a one week interval (September 20th and 26th), showed COTS density and abundance decreased and the seastars appeared to be migrating slowly in a northerly direction and into shallower waters.
- Coral tissue loss resulting from biological interactions such as COTS predation also has the ability to
- influence benthic community structure and coral health by making substrate available for algal colonization and may have a large, but underestimated, influence on coral health, benthic community structure and percent live coral cover along West Hawai'i.
- Only a single predator of COTS, *Charonia tritonis* (Triton's Trumpet), was observed at the survey site.
- Continued monitoring of COTS populations and immediate protection of their predators, *C. tritonis* and *Cassis cornuta* (Triton's Trumpet and Horned Helmet) is highly recommended throughout the State of Hawai'i.
- Three species of monitored urchins have been increasing on West Hawai'i reefs with the collector urchin, *T. gratilla* exhibiting the greatest increase. This increase is not related to an increase in benthic algae as a food supply.

East Hawai'i

- Abundance of fishes is significantly greater at both Waiopae MLCD and Waiopae open sites than at Richardson's Ocean Center (ROC). Species richness is higher in the MLCD as compared to ROC. The MLCD and ROC sites have the highest similarity in fish communities, and the OPEN and ROC communities have the lowest similarity.
- Over the 12 years of surveying of fishes at Waiopae and ROC, there appears to have been a slight increase in fishes observed between 1999 and 2006, followed by a three-year decline. No net increase in fish abundance has been observed at Waiopae MLCD since its establishment in 2003.

CONTRIBUTORS

Laura Livnat, Ross Martin, Kara Osada-D'Avella, Rich Osada, , Kosta Stamoulis, Brian Tissot, Ivor Williams, Jill Zamzow, Greta Aeby Megan Lamson, Lindsey Kramer, Rebecca Most, Anthony Spitzack, Chad Wiggins, Brent Carman.

Hawai'i Island Surveys

Benthic Monitoring

Methods

Benthic surveys were initially conducted in West Hawai'i in 1999 and then again in 2003. More recently, surveys were conducted at 26 monitoring sites in 2007 and 2011. The images used for analysis in 1999 were captured by digital video. The resolution of the video images was very poor however compared to the subsequent surveys which used much higher resolution digital still images (Olympus 5060 in 2003, Olympus 7070 in 2007 and Olympus E-PL1 in 2011). Specifically, octocoral was not detectable in the 1999 video capture images, nor was it possible to distinguish live finger coral from dead finger coral. It was therefore determined that it was not valid to compare data taken with these two different techniques.

To obtain images of consistent size and quality, a 75cm clear Plexiglas® spacer rod is attached to the underwater housing and used as a guide to steady the camera at a fixed height (0.75m) above the benthos. A white balance feature was used to compensate for loss of red light at depth, giving the images a more natural appearance without artificial lighting. Four transects 25m in length were photographed at each site. Images were taken at 1m intervals from a standard height of 0.75cm starting at the 0 point and ending at the 25m mark, producing 26 images per transect.

Images were analyzed using the Coral Point Count with Excel extensions software program (CPCe Kohler and Gill 2006). Data was pooled by transect. The resulting configuration was 4 transects per site, 26 frames per transect, 20 stratified random points per image (4 rows, 5 columns), 520 individual data points per transect, and 2080 points per site. Proportion of each benthic category was determined for each image and percent cover was calculated for each transect, Total percent cover was obtained by calculating the mean percent cover of the 4 transects.

Results

Complete benthic data for the 2003, 2007 and 2011 surveys, presented as percent coverage, are contained in Appendices A – H. Comparisons of total coral cover (paired two-sample T tests) were performed on the percent total coral cover mean values for individual transects (Table 1).

Between Lapakahi, the northernmost site, and Keahole Point, a distance of approximately 37 coastal miles, there are 9 survey sites. One site, Unualoha, was added in 2007 and therefore no comparative data is available. Of the 8 “northern” sites (north of Keāhole Point) that were compared, 6 showed statistically significant declines in total coral cover between 2003 and 2007. Lapakahi, Kamilo Gulch and Waiaka'ilio Bay (the three northernmost sites), Keawaiki, 'Anaeho'omalū and Ka'upulehu all declined significantly. Only Puakō and Makalawena showed no significant change (Figure 37).

A severe storm with large swells caused extensive coral damage along the West Hawai'i coast north of Keāhole Point in January 2004. This damage was noted during surveys soon after the storms occurred. The declines at Kamilo Gulch and Waiaka'ilio Bay may also have been influenced by a major sediment runoff event caused by heavy rainfall in October 2006. A reconnaissance was conducted offshore of several intermittent streams near these sites soon after the event. Thick layers of sediment covering large amounts of coral were observed and sediment was recorded at water depths of 90 feet. Numerous dead coral were observed during subsequent reconnaissance.

South of Keāhole Point 15 sites were compared. Three sites, Wawaloli Beach, Papawai Bay and South Oneo Bay showed statistically significant increases in total coral cover between 2003 and 2007. All other sites showed no change.

Table 1. Percent coral cover at West Hawai'i sites in 2003, 2007, 2011

Site (N to S)	2003	2007	2011	Δ (2007/2011)	P =	Δ (2003/2011)	P =	
Lapakahi	19.50%	11.37%	11.78%	0.41	0.794	-7.69	0.032	Overall Decline, no change from 2007 to 2011
Kamilo	49.50%	38.19%	28.96%	-9.23	0.000	-20.27	0.004	Overall Decline
Waiaka'ilio Bay	54.40%	42.50%	38.78%	-3.72	0.089	-15.13	0.054	Overall Decline, no change from 2007 to 2011
Puakō	49.09%	47.83%	34.21%	-13.63	0.001	-14.88	0.002	Overall Decline mainly between 2007/2011
'Anaeho'omalu	40.58%	31.47%	28.43%	-3.04	0.147	-12.15	0.005	Overall Decline, no change from 2007 to 2011
Keawaiki	29.66%	16.73%	18.68%	1.95	0.220	-10.98	0.031	Overall Decline, no change from 2007 to 2011
Ka'ūpūlehu	40.71%	31.15%	27.05%	-4.09	0.171	-13.66	0.030	Overall Decline, no change from 2007 to 2011
Makalawena	44.88%	47.57%	47.63%	0.06	0.992	2.76	0.489	No Change
Unualoha	N/A	36.82%	36.51%	-0.31	0.873	N/A	N/A	No Change (only measured from 2007/2011)
Wawaloli	37.21%	37.51%	42.26%	4.74	0.061	5.05	0.140	No overall change, increasing from 2007/2011
Wawaloli Beach	37.93%	42.25%	44.45%	2.20	0.479	6.52	0.187	No Change
Honokōhau	43.22%	48.54%	48.32%	-0.22	0.940	5.10	0.437	No Change
Papawai	32.31%	38.31%	41.05%	2.75	0.506	8.84	0.173	No Change
Old Kona Airport	N/A	53.16%	51.19%	-1.97	0.570	N/A	N/A	No Change (only measured from 2007/2011)
S. Oneo Bay	56.09%	61.86%	46.55%	-15.31	0.019	-9.54	0.054	Overall Decline mainly between 2007/2011
N. Keauhou	31.92%	31.28%	28.00%	-3.28	0.134	-3.92	0.165	No Change
Kualanui	52.81%	62.35%	59.78%	-2.57	0.358	6.97	0.124	No Change
Red Hill	30.68%	33.22%	35.26%	2.04	0.470	4.58	0.148	No Change
Keopuka	15.98%	15.59%	14.44%	-1.15	0.600	-1.54	0.559	No Change
Kealakekua Bay	27.10%	28.64%	23.11%	-5.53	0.288	-3.99	0.219	No Change
Ke'ei	31.20%	28.67%	26.70%	-1.96	0.543	-4.50	0.379	No Change
Kalahiki	36.53%	39.62%	38.94%	-0.68	0.720	2.41	0.026	No Change
Ho'okena (Auau)	28.18%	28.44%	29.98%	1.54	0.671	1.80	0.109	No Change
Miloli'i (Omaka'a)	29.76%	27.08%	32.94%	5.86	0.052	3.18	0.414	Increase from 2007/2011
Manukā	30.35%	33.17%	33.36%	0.19	0.961	3.01	0.689	No Change




Figure 1. Comparison among survey years of percent coral cover across West Hawai'i monitoring sites

Puakō

The situation at Puakō merits special attention. For the 2003 and 2007 surveys, percent coral cover at Puakō was statistically unchanged at 49.9% and 47.8% respectively. In 2011 there was a significant decrease in coral cover to 34.2% (Figure 2).




Figure 2. Same location over time showing significant loss of *Porites lobata* (Δ -13.6% between survey years)

Minton, et al. (2012) reports a severe decline in coral cover from 80% in the mid 1970's to 32% in 2010. (DAR 2011 analysis showed 34.1% coral cover in 2011). The report theorizes a number of causes primarily due to human impacts (overfishing and local land based pollution). Plans for development in the area call for 4,000 new homes to be built. With this staggering increase the potential for catastrophic anthropomorphic impacts to the reefs in this area should not be ignored. The South Kohala Conservation Action Plan developed by a multi-agency team addresses many of the problems affecting this shoreline. Coastal development will inevitably increase in West Hawai'i. This plan addresses one relatively small but important area of the coast. More conservation plans of this nature should be considered before irrevocable impacts occur.

In addition, surveys conducted in 2010 and 2011 as part of a coral health monitoring program conducted by Courtney S. Couch (Cornell University Ph.D candidate) in collaboration with DAR showed algal overgrowth and the resulting coral tissue mortality occurred at all eight DAR sites surveyed, including Puakō (WHAP Site 4). In the majority of cases of active algal overgrowth, the red filamentous algae *Corallophila huysmansii* was the primary contributor to coral tissue mortality (Couch et al. in prep) and therefore may be also be contributing to the significant loss of coral cover at Puakō.

Octocoral Distribution

Benthic surveys revealed a most interesting distribution of one or more species of octocorals centered on the urbanized areas of Kailua-Kona (Figure 3). The Bishop Museum checklist (<http://www2.bishopmuseum.org/HBS/invert/results.asp>) lists 11 species of shallow water octocorals occurring in Hawai'i. At least one of the species in question appears to be the blue octocoral *Sarcothelia (Anthelia) edmondsoni* -Figure 38) although the taxonomy of the group is somewhat confused. The original taxonomic description for *S. edmondsoni* is actually a brown morph common in calm lagoons on the windward side. The blue morph is more abundant in fore reef areas with heavy wave surge and is most likely a separate species. Both varieties have long histories in Hawai'i and are presumably native and/or endemic (S. Kahng, pers. comm.)

The apparent concentration of *Sarcothelia edmondsoni* (Figure 39) in the vicinity of Honokōhau Harbor and Kailua Bay (Figure #) may suggest anthropogenic influence on the distribution of octocoral in West Hawai'i. Published studies have suggested that octocorals may be indicators of pollution (Baker and Webster 2010, Hernandez-Munoz et al. 2008). With the planned increase in development in these areas and the possible associated rise in point source pollution further investigation into octocoral distribution and its potential as a pollution indicator is suggested.

An analysis of octocoral percent cover changes between 2003 and 2007 showed no statistically significant change in all but one of the West Hawai'i sites where octocoral has been recorded. Papawai Bay declined from 18.2% to 10.9% ($P=.018$). The next benthic surveys to be conducted in early 2011 will permit octocoral percent cover comparisons of the 6 sites that were added after 2003.



Figure 3. *Sarcothelia edmondsoni* (left) and another unidentified octocoral found on West Hawai'i reefs




Figure 4. Comparison between survey years of percent cover of octocoral across West Hawai'i monitoring sites

Coral Disease

Methodology

Coral disease surveys were conducted at 28 West Hawai'i sites, and at Okoe and Hōnaunau Bays. Surveys were conducted from March to July 2010 by four survey divers: Courtney Couch (Cornell University), Camille Barnett (DAR), Kara Osada-D'Avella (DAR), and Linda B. Preskitt (DAR). Two permanent transects were surveyed at each site.

Field surveys

An area of 1 x 25 meters was surveyed for coral disease along each transect. Larger areas were surveyed at sites with low occurrences of disease, while (due to time constraints) the full 25 m² was not surveyed at several sites with high disease frequency. Disease assessment included all corals within the survey area inspected for signs of trematodiasis, growth anomalies, tissue loss syndrome, multifocal tissue loss, hypermycosis, and other progressive conditions. When disease was present, colony size and species were recorded along with the number, size, shape and color of the lesion(s) observed. All diseased colonies were photographed and described, excluding colonies with only *Porites* trematodiasis. In addition, 1-2cm fragments from diseased

coral colonies were sampled for histological analyses, helping to further differentiate between tissue loss and biological interactions (e.g. predation).

Colony assessment

Colony counts were conducted in conjunction with coral disease surveys. For each transect line, a 1 x 10 meter area was surveyed with the aid of a 1m square quadrat. Each coral colony within the survey area was recorded to species level and assigned to one of seven size classes; 0-5cm, 5.1-10cm, 10.1-20cm, 21.1-40cm, 41.1-80cm, 81.1-160cm and >160cm.

Calculations and Analyses

We calculated mean colony density (colonies/m²) for each site by averaging the number of colonies of each genus on both transects and dividing by the average area surveyed for each site. Mean colony density was then multiplied by the area surveyed for disease to obtain estimated number of colonies. At each site we calculated total estimated disease prevalence for each disease as follows: (total no. cases of a specific disease for the genus) ÷ (estimated number of colonies for the genus). Total disease prevalence for each site was calculated using the method described above using total number of colonies and total number of diseased cases for each site (all genera combined).

In 2007, Dr. Greta Aeby and Steve Cotton (DAR) conducted initial coral disease surveys at 10 WHAP sites (Appendix Table A) (DAR 2007). This dataset was compared with data collected in 2010 to assess changes in coral disease frequencies. Prevalence (% of diseased colonies per site) data between the two surveys were not comparable due to substantial differences in colony counts between 2007 and 2010, with significantly more small colonies (colonies <10cm) counted in 2010 than 2007. This difference was believed to be due to observer changes rather than biological changes. Therefore, data were compared using disease abundance per m² rather than disease prevalence.

Coral disease prevalence data were non-parametric; therefore Spearman rank correlation analyses were employed. Paired t-tests were used for comparisons of disease per m² between 2007 and 2010 surveys at ten WHAP sites (JMP® v8.0.2.2, ©2009 SAS Institute Inc.)

Results

Coral disease by size class

Coral diseases were observed across all colony size classes, with the greatest percentages of disease cases occurring in the larger size categories (Figure 40). Coral colonies less < 5cm accounted for 18% of total colonies (18.3% of *Porites* spp.) recorded in count surveys, yet accounted for only 1% of the total cases of diseased colonies (1.1 % of *Porites* spp.). These findings imply West Hawaii's small corals (<5 cm) are less susceptible to disease than the larger and subsequently older colonies. Linear growth rates of coral colonies are both species and size specific, and are affected by a suite of environmental factors such as depth, temperature, light irradiation and latitude.




Figure 5. Size structure of coral colonies recorded in West Hawai'i during surveys conducted in 2010. Percentages reflect % of diseased colonies recorded within each size class

Therefore it is difficult to age colonies based on size. Given an overall slow growth rate (ranging from 7.4 – 16.7 mm/yr.) of the dominant reef builder *Porites* and the relative contribution of gametes that large colonies provide, it is important to continue monitoring coral disease prevalence as they may have long-term effects on coral populations and community structure (Rodgers & Cox 2003; Forsman et al. 2006, Richmond 1987; Grigg & Maragos 1974, Lough & Barnes 2000).

Why more diseased colonies than total colonies recorded?

Larger colonies tended to occur near the end of survey lines, therefore the number of diseased colonies are greater than total colonies counted due to the methodology employed; coral colonies were counted and sized for the first 10m of each transect, while disease assessments were made along the full 25m line.

Coral community structure

Percent coral cover varied across surveyed sites, ranging from 11.5% at Site 1 (Lapakahi) to 62.0% at Site 14 (South Oneo Bay) (Appendix Table). Within all monitoring sites, *Poritids* were the most abundant corals, while densities of other coral genera were variable across sites. Coral colony density was not significantly related to

percent coral cover (Spearman $\rho = -0.2626$, $p > 0.1$). Rather, high coral density reflects an abundance of small colonies (Figure 40).


Spearman rank correlations revealed significant negative relationships between overall colony density and total disease prevalence ($\rho = -0.5276$, $p = 0.0033$). However, total disease prevalence was positively related to percent coral cover ($\rho = 0.4291$, $p = 0.0202$). In other words, higher disease prevalence was observed on reefs with high coral cover and lower colony density, which is likely due to the increase in disease susceptibility with colony size.

When relationships were analyzed by genus, *Porites* followed the same trend as described above. *Porites* growth anomalies and *Porites* tissue loss syndrome were positively correlated with percent cover of *Porites* sp. ($\rho = 0.4444$, $p = 0.0178$ and $\rho = 0.3804$, $p = 0.0458$) and negatively related to Poritid density ($\rho = -0.7200$, $p < 0.001$ and $\rho = -0.5600$, $p = 0.0016$). Frequency of *Porites* diseases may be attributed to the dominance of Poritid corals in West Hawai'i reef communities possibly allowing the spread of pathogens or creating a susceptibility to disease within the genus.

Coral diseases in West Hawai'i

At 30 sites surveyed in West Hawai'i, the following diseases were recorded within each specified genus: growth anomalies (GA) of Poritids and Montiporids, *Porites* trematodiasis (TRE), tissue loss syndrome (TLS) within *Porites* and *Pocillopora*, *Porites* multifocal tissue loss (MFTL), and hypermycosis (HYP) of *Pavona* (Figure 41).

The above diseases have been previously described (Coral Disease Working Group 2007, Williams et al. 2010), however we observed a number of cases of a distinct type of tissue loss in *Pocillopora meandrina*. The lesion was characterized by progressive tissue loss from one side of the colony with old algae-covered skeleton grading into recently denuded skeleton to sloughing and into apparently healthy tissue. The tissue loss appears to originate and progress from the base of each branch, with a clear band of freshly denuded skeleton at the lesion margin. We also recorded cases of possible *Pocillopora* senescence. This condition is common along West (C. Couch pers. obs.) and East Hawai'i (B. Vargas-Angel pers. comm.) In most cases colony death originates on one side of the colony and progresses across the colony. Algal covered skeleton is adjacent to pale/bleached tissue, which grades into "normally" pigmented tissue. Samples sent to United States Geological Survey (USGS) Biological Resources Division for analyses revealed atrophy, appearing to be a senescence reaction (or progressive death of the colony, perhaps due to age) (Figure 41).






Figure 7. Relative abundance of coral diseases recorded for DAR monitoring sites in West Hawai'i during surveys conducted in 2010.

Although *Porites* growth anomalies were found at all but two sites (Sites 6, Keawaiki and Site 97, Unualoha Pt.), mean prevalence across all sites was low (1.83 ± 2.15 %), ranging from 0.02 % at Site 10 (Wawaloli Beach) to 7.81% at Site 11 (Honokōhau) (Figure 43).

Porites trematodiasis, the second most common disease, was found at all but the following four sites: Site 2 (Kamilo Gulch), Site 3 (Waiaka'ilio), Site 18 (Keopuka), and Site 21 (Kalahiki Beach). Mean prevalence across all sites was low (1.71 ± 2.17 %), ranging from 0.05 % at Site 1 (Lapakahi) to 9.03% at Site 8 (Makalawena) (Figure 43). *Porites* tissue loss syndrome occurred at all but the following sites: Site 1 (Lapakahi), Site 6 (Keawaiki), Site 7 (Ka'ūpūlehu), Site 97 (Unualoha Pt.), and Site 17 (Red Hill). Mean prevalence across all sites was low (0.21 ± 0.18 %), ranging from 0.02 at Site 10 (Wawaloli Beach) to 0.65 % at Site 23 (Omaka'a Bay) (Figure 43).




Figure 8. Prevalence of Poritid diseases at each West Hawai'i site surveyed in 2010

Although possible senescence reaction of *Pocillopora meandrina* appears commonly in West Hawai'i (see section entitled Diseases in West Hawai'i), this condition was observed at only two sites: 97 (Unualoha Pt.) and 22 (Ho'okena). This infrequent documentation of cases is likely attributed to the low number of *Pocilloporids* occurring at DAR monitoring sites, as *P. meandrina* accounts for an average of 0.83% of total coral cover at WHAP sites

Spatial Patterns

Anthropogenic impacts such as coastal pollution are hypothesized to result in physiological stress and altered host-pathogen interactions, leading to changes in coral health and coral reef community structure (Harvell et al. 2007). While the mechanisms underlying the link between coral disease and water quality are poorly understood, diseases such as growth anomalies have been positively associated with high human use and impaired water quality in the Pacific (Yamashiro et al. 2000, Kaczmarzsky 2009, Aeby et al. 2011).

Due to Hawai'i's highly porous basaltic rock, terrestrial inputs are transported rapidly through submarine groundwater (Knee et al. 2010). Data collected by Johnson (2008)

documented areas with submarine groundwater discharge (SGD) “plumes” between Kawaihae and Hōnaunau. Disease prevalence at DAR monitoring sites was analyzed in relation to these SGD plumes (data were available for a total of 14 monitoring sites within the region documented).

Overall disease prevalence and prevalence of *Porites* growth anomalies were positively correlated with total estimated size of SGD plumes (total prevalence $r = 0.460$, $p = 0.098$, *Porites* GA $r = 0.586$, $p = 0.028$) and number of SGD plumes (total prevalence $r = 0.612$, $p = 0.020$, *Porites* GA $r = 0.744$, $p = 0.002$) located within the vicinity of each site (<1.5 km). These results show high nutrient loading may be affecting West Hawai'i's coral health.

Additionally, sites surveyed in West Hawai'i show a significant negative relationship between disease prevalence and distance from harbors/boat ramps (overall disease prevalence: $r = -0.402$, $p = 0.028$) (Figure 44). The most frequently occurring diseases, *Porites* growth anomalies and *Porites* tissue loss syndrome showed decreased prevalence with greater distance to these usage areas (*Porites* GA $r = -0.701$, $p = 0.0001$, *Porites* TLS $r = -0.658$, $p = 0.0001$). Similar to previous findings, the distribution of *Porites* trematodiasis, a disease known to be transmitted by fishes, particularly corallivores, was not associated with these locations (Aeby 2007).




Figure 9. Disease prevalence in relation to site distance from harbors/boat ramps in West Hawai'i for overall disease prevalence ($r = -0.402$, $p = 0.028$), *Porites* growth anomalies (GA) ($r = -0.658$, $p = 0.000$) and *Porites* tissue loss syndrome (TLS) ($r = -0.701$, $p = 0.000$)

Prior studies have also found relationships between abundances of reef fish and prevalence of particular coral diseases. Various fishes are known to impact corals directly (such as the grazing of parrotfish) as well as transmit diseases (such as corallivorous butterflyfish) (Williams et al. 2010). Aeby et al. 1998 also found the highest

trematodiasis at sites with intermediate percent coral cover. Using fish abundance data from WHAP surveys, sites were compared for Poritid disease prevalence to corallivorous butterflyfish and parrotfish abundances. However, no statistically significant relationships were found between these fish groups and coral disease prevalence for West Hawai'i's reefs.

Temporal comparisons

Comparisons of disease density (instances per square meter) between 2007 data and 2010 revealed no significant changes in disease densities between survey years ($t = -1.46$, $p = 0.18$). Though changes were not significant, *Porites* trematodiasis slightly increased at most sites (Figure 45).




Figure 10. Comparison between survey years (2007 vs. 2010) of diseased colony densities for three types of Poritid conditions at 10 sites in West Hawai'i

Instances of *Porites* growth anomalies and *Porites* tissue loss syndrome increased (though not significantly) at four sites: Sites 4 (Puakō), 5 (Mauna Lani), 11 (Honokōhau) and 15 (Keauhou). Each of these sites is located in close proximity to harbors and boat ramps. As described in the previous section, diseases have been positively associated with high human use.


Although no significant change in disease frequency was found, the change in presence or absence of two diseases was noted. *Montipora* white syndrome was not recorded in surveys in 2010, though one case was recorded in 2007. *Pocillopora* tissue loss (including senescence reaction) and *Pavona varians* hypermycosis were not recorded in 2007 surveys, but occurred at multiple sites in 2010. For *Pavona varians* hypermycosis, this includes some sites previously surveyed (Table 8).

Temperature data


Hobo[®] temperature loggers (Onset Computer Corporation) were initially deployed at all West Hawai'i Fish Replenishment Area (FRA) sites (Figure 46). They were attached via cable tie to a coral head in the immediate vicinity of the center transect pin. Due to various circumstances including loss and flooding (i.e. multiple Hobo[®] Water Tem Pro units) a complete temperature record over the last decade is not available for any site. Fortunately fairly comprehensive temperature data exists for several West Hawai'i sites including a northerly site (Waiaka'ilio), a southerly site (Miloli'i) and a central site (Ke'ei) (Figure 47).

Examination of the temperature data reveals a marked similarity in water temperatures along coastal sites separated by considerable distances. From 1999 to 2005 there was a clear trend of increasing water temperatures along the West Hawai'i coastline. Over this 6 year period water temperatures increased by 1.8-2.7°F. For comparison, surface water temperature records at Koko Head, O'ahu indicated an increase of 1.4°F over a 50 year period (NMFS + IGLOSS corrected data provided by Paul Jokiel). Trend analysis suggested that if West Hawai'i water temperatures continued to increase unabated, the lethal thermal limit for corals (i.e. 30 day exposure to mean water temperature of 29.6°C) would likely be reached within a decade. The good news is that waters have not continued to increase and actually have decreased over the past several years.


The most recent El Niño event to occur began in June 2009, peaked in November and December of the same year and waned in March 2010. It was effectively over by June 2010 (Jet Propulsion Laboratory web site). Although El Niño periods are characterized by warmer than usual equatorial waters, West Hawai'i coastal waters were only marginally warmer than the preceding two years. Mean water temperatures for the four month period of Oct 09-Jan10 was 78.9°F which was only 0.4 - 0.5°F warmer than the previous periods (Oct-07-Jan 08 = 78.5°F; Oct 08-Jan 09 = 78.6°F). Examination of the temperature records also shows that water temperatures in several of the previous years (e.g. 2004/2005) were generally higher than during the recent El Niño event.



Waiaka'ilio FRA



Ke'ei FRA



Miloli'i FRA

Figure 11. Temperature (°F) records for three FRAs

Fish Surveys

Although the DAR fish survey protocol for West Hawai'i was initially designed to focus primarily on species which are the principal targets of the aquarium fishery it has proven to be a highly useful methodology for general coral reef monitoring and has been adopted by DAR for monitoring on other islands. It's important to note that all fishes are censused, whether they're aquarium species or not. While the protocol is particularly effective for assessing recruits, smaller site-oriented species and those not wary of divers; it also provides highly useful information on other groups including predators, invertebrates and "food" fishes. The specifics of the methodology are detailed in the O'ahu section (pg. 13).

DAR monitored 23 sites in West Hawai'i (Figure 2) bi-monthly, for a total of six surveys per year (five in 2000 due to logistic problems) until Jan. 2005 when the project was revamped at which time surveys became quarterly.




Figure 12. West Hawai'i monitoring sites

These fixed transect surveys are noted as 'small scale' surveys in Table 9. Similar monitoring has also been conducted at three sites in East Hawai'i although on a less systematic schedule.

In addition to the transect surveys, a 10 minute 'free-swim' survey is also conducted by two divers in the areas surrounding the fixed transects. The purpose of this survey is to increase the ability to census uncommon or rare species and species of particular ecological interest such as cleaner wrasse (*Labroides phthiophagus*), ta'ape (*Lutjanus kasmira*), roi (*Cephalopholis argus*), crown-of-thorns (*Acanthaster planci*) and all species of terminal phase parrotfishes. Recording of species during the timed free-swim survey that were not observed on the transect surveys augments a site-specific species list.

In order to obtain better data on fish species that are heavily harvested and in demand for both subsistence, recreational and commercial food fisheries (i.e. 'resource fish') an enhanced monitoring protocol was newly implemented in 2005 at all new survey sites and at a number of existing monitoring sites (Table 9). 'Resource fish' are surveyed by a pair of divers swimming in parallel (10m apart), following a depth contour, for a five minute period. Each diver records all 'resource fishes' (main fishery target species) >15cm within a 5m wide belt. Rare, skittish or uncommon fishes such as sharks, rays or carangids which are observed any time throughout the survey dive are noted. Starting points for this survey are based on existing center pin site coordinates. End points are delimited by a diver deploying a surface float at the completion of the 5 minute survey. Sites which include all three types of monitoring are termed "Integrated" (Table 9).

Table 3. West Hawai'i monitoring sites with corresponding coordinates, status and survey type (INT=Integrated monitoring, SS=Small scale, RF=Resource fish only)

Site	District	Latitude	Longitude	Mean Depth (m)	Status	Type
Lapakahi	N. Kohala	20.1600000	-155.9001833	12.1	MLCD*	INT
Kamilo Gulch	N. Kohala	20.0810167	-155.8680833	12.8	Open	SS
Waiaka'ilio	N. Kohala	20.0739167	-155.8645167	13.4	FRA	INT
Puakō	S. Kohala	19.9698833	-155.8488000	9.2	FMA	INT
'Anaeho'omalū Bay	S. Kohala	19.9527500	-155.8661667	10.0	FRA	INT
Keawaiki	N. Kona	19.8911167	-155.9100667	13.3	Open	SS
Ka'ūpūlehu	N. Kona	19.8439500	-155.9809667	11.4	Open	SS
Makalawena	N. Kona	19.7965000	-156.0328833	10.2	FMA	INT
Ho'ona / Unualoha Pt.	N. Kona	19.7425100	-156.0557500	12.4	Open	INT
Wawaloli Beach	N. Kona	19.7088833	-156.0494951	9.8	Open	SS
Wawaloli	N. Kona	19.7000100	-156.0499100	13.6	Open	SS
Kaloko-Honokōhau	N. Kona	19.6709833	-156.0303333	13.1	FRA	INT
Papawai	N. Kona	19.6472500	-156.0229833	10.4	FMA	SS
Old Kona Airport	N. Kona	19.6421200	-156.0121000	12.2	MLCD	INT
S. Oneo Bay	N. Kona	19.6312000	-155.9930000	12.0	FRA	SS
Keauhou	N. Kona	19.5683833	-155.9693500	12.0	FRA	INT
Kualanui Pt. (Red Hill)	N. Kona	19.5482667	-155.9623000	11.3	Open	SS

Red Hill	S. Kona	19.5052833	-155.9528833	13.9	FMA	INT
Keopuka	S. Kona	19.4829167	-155.9460000	10.3	Open	SS
Kealakekua Bay	S. Kona	19.4793000	-155.9327833	8.0	MLCD	INT
Ke'ei	S. Kona	19.4628167	-155.9268000	11.5	FRA	INT
Ho'okena (Kalahiki)	S. Kona	19.3691500	-155.8974000	11.1	FRA	SS
Ho'okena (Auau)	S. Kona	19.2978833	-155.8898833	13.6	Open	SS
Miloli'i/Honomalino	S. Kona	19.1673000	-155.9132500	12.3	FRA	INT
Okoe Bay	Ka'u	19.6421200	-156.0121000	16.5	FRA	RF
Manukā	Ka'u	19.0767167	-155.9039667	12.0	Open	SS

Shallow Water Resource Fish Surveys

Shallow water resource fish surveys collect data on the abundance of resource (desired) fish species in shallow water habitats where they are typically most abundant during the day in West Hawai'i. These surveys were designed to be comparable with our standard




Figure 13. Map showing locations of shallow water resource fish surveys and laynet prohibited areas

resource fish surveys occurring in mid-depth habitats, and thus the methodology is very similar. As with the other resource fish surveys, distance covered is measured for every survey so that data can be analyzed on a per unit area basis. Initially 72 sites were

selected evenly distributed along the coastline in 2-6m of water between our northern and southernmost permanent study sites (Figure 48). Using a GIS (ArcGIS 9.2), the 72 points were overlaid on a NOAA habitat map for the purpose of adjusting any sites that did not fall on hard-bottom habitat. Direction taken for the survey was predetermined when habitat was an issue. Otherwise survey direction (north or south) from the start point was determined in the field. Each site is surveyed only once.

The survey consists of a timed 10min swim along the coastline with divers being careful to remain in the target depth of 2-6m. When the survey is finished, the boat captain records an end point so that the distance covered can be later calculated. The dive team consists of two divers both surveying a single 5m wide belt. One diver is counting surgeonfish, goatfish, and introduced species above 15cm except for *Acanthurus achilles* and *A. triostegus* for which individuals above 10cm are recorded. The other diver counts parrotfish, wrasses, other resource fish, and selected rare butterflyfish of interest. Large predatory fish appearing off transect are also recorded.

Adult Yellow Tang surveys

To supplement data from the long-term monitoring program and to investigate the possibility of 'spillover' of adult fish from existing protected areas, we survey adult yellow tang populations in their prime daytime habitat, i.e. the deep edge of the shallow pavement zone around 3 to 6 m deep. Along the West Hawai'i coast, shallow pavement areas generally have a distinct deep boundary where the main reef slope begins and where coral cover increases rapidly, and therefore the target habitat zone for our surveys was mostly well defined. Recognizing that adult yellow tang have highly clumped distributions, we developed a survey approach which allows divers to count yellow tang over long transects running approximately parallel to shore through the prime adult habitat.

There are 4 AYT sites within FRAs, 4 within long-term protected areas (LTP); and 8 in open, i.e. fished, areas. As adults have daily movements between diel and night time areas of up to at least 800 m we assumed that there could be spillover across protected area boundaries over at least that scale. We therefore established 4 open sites as 'boundary' sites, centered < 1 km from the nearest protected area boundary, and 4 as 'open' sites with mid-points > 2 km from the nearest boundary. Each area was surveyed 5 times in 2006 and 6 times in 2010 and analysis of the latest data will commence in the new year. The survey technique and initial findings of significant spillover of yellow tang from protected to open areas is contained in Williams et al. 2009.

Depth Stratified Random Surveys

In response to a long standing conflict between aquarium fish collectors and the local community at Ka'ohē (Pebble Beach), South Kona, a DLNR community advisory group, the West Hawai'i Fisheries Council (WHFC) recently recommended that the area at Ka'ohē be closed to aquarium collecting. To maintain the existing balance of open and closed areas the WHFC also recommended that a similarly sized protected area be opened at to collecting Keauhou which is presently in an FRA. Considerable disagreement ensued however surrounding the nature and abundance of the resources within the proposed open area so DAR embarked on an effort to accurately assess the populations of a number of species of interest. 72 random, depth stratified, surveys were conducted at Keauhou (Figure 49) in July 2008 to derive area population

estimates. Survey methodology closely follows the methodology described above for 25m fixed transects but with two rather than four 25mX4m transects at each random point. The Keauhou survey was repeated in August 2010 and similar surveys have been conducted at Ka'upulehu (August 2009) and Red Hill (April 2009).




Figure 14. Map showing the locations of Keauhou stratified random fish population survey sites. The stratified depth zones are as follows: DP=24-30m, MD=18-24m, MI=9-18m, SH=3-9m

Retrospective Surveys

Several long-term retrospective surveys, primarily directed at fish populations, are being conducted at 3 West Hawai'i sites. The sites and the date of the initiation of the original surveys are as follows: Puako, South Kohala (1979), Ke'ei, South Kona (1978) and Honaunau, South Kona (1975). So that new data is comparable with historical data, the same transect locations and survey methodologies are employed as in the original studies. Methods vary by locations, but all are based on standard dimension belts or search areas. Additional benthic data are also being collected. This work is presently under analysis.

Results

Fish Surveys

West Hawai'i

Fishes on West Hawai'i reefs may be regarded as falling into three groups based upon human utilization. Resource or 'food' fish such as jacks (Carangidae), goatfishes (mullidae) and parrotfishes (Scaridae) are those targeted for food by recreational and commercial fishers. Aquarium fish are those which are harvested, usually in the smaller size classes, by commercial aquarium collectors although there are some species which fall into both groups (e.g. kole, *Ctenochaetus strigosus* and Achilles tang, *Acanthurus achilles*) for the present study these are classified solely as aquarium fishes. The third group ('other') is species which are harvested neither for food nor for aquaria.

The overall number of 'other' fishes, those which are not substantially harvested for either food or for the aquarium trade, did not change significantly at West Hawai'i sites over the last 11 years although individual species within this group may have. In contrast, the abundance of both aquarium and food fishes increased significantly over the same time period (Figure 15). For aquarium fishes it is clear that a substantial part of the increase in overall numbers is due to the implementation in 2000 of a network of





Figure 15. Overall temporal trend in mean fish density of three major fish utilization groups at West Hawai'i sites. Aquarium Fishes represents top 20 collected species. Trend line represents LOESS (locally weighted polynomial regression) smoothing procedure applied to data. Closed triangle = $p < 0.05$ (Spearman rank test)

Fish Replenishment Areas (FRAs) along the West Hawai'i coast. The aquarium fishery in Hawai'i is economically the largest inshore fishery in the state and certainly the most conflict filled. The management importance of comprehensive and extensive monitoring such as has been underway in West Hawai'i for over a decade cannot be underestimated when addressing the issue of this highly controversial fishery. In depth analysis of aquarium collecting impacts is contained in a later section (pg. 54).

Examination of the temporal trends of some of the most common reef fish families indicates that acanthurids have been increasing over the past eleven years while labrids have decreased (Figure 16). Overall, chaetodontids and pomacentrids have been relatively stable although some species within the family have either increased or declined.






Figure 16. Temporal trend in mean fish density for four families on West Hawai'i sites. Trend line represents LOESS (locally weighted polynomial regression) smoothing procedure applied to data. Open triangle = $p < 0.1$, Closed triangle = $p < 0.05$ (Spearman rank test)

As noted, the species within each family can vary substantially in temporal trends as exemplified by the wrasses where three species increased, two decreased and three remained stable. The reasons for such differences are not clear but it does appear that with only a single exception (*Stethojulis balteata*) most wrasses are in a period of decreasing abundance, undoubtedly influenced by low levels of recent recruitment. Among the acanthurids the two species exhibiting the most substantial increases (*Zebrasoma flavescens* and *Ctenochaetus strigosus*) are also the most heavily collected aquarium species comprising approximately 91% of the total catch. It is also apparent however that a number of less abundant aquarium-targeted species such as the moorish Idol (*Zanclus cornutus*) and lei triggerfish (*Sufflamen bursa*) have not responded to the increase in protected areas and have actually decreased in West Hawai'i since 1999 (Figure 17).




Figure 17. Temporal trend in mean fish density for various fish species of interest at West Hawai'i sites. Trend line and triangles as above




Figure 18. Overall change in herbivore biomass in FRA, MPA and open areas 2003-2012

Introduced species

Ta'ape

From their initial introductions, ta'ape have clearly undergone an expansive period of population growth. Ta'ape were only introduced to the island of O'ahu but have subsequently spread widely throughout the islands of the archipelago. Based on free swim site surveys there was a trend for increasing numbers from 1999 to 2004 followed by a subsequent of unknown cause.

Transect data reflects overall low abundance of this species in the reef areas of the study sites (2010-2012 mean = 0.14/100m²). Similarly ta'ape are rarely found in the shallower water where resource fish surveys are conducted (mean = xx/100m²). While Ta'ape are numerous in some locales usually along drop-offs and deeper reef areas, their distribution is highly patchy (characteristic of a schooling species) and they are not at all abundant in many reef areas in West Hawai'i. Similar to West Hawai'i, at some shallow reef locations such as in Kāne'ohe Bay, ta'ape numbers also appear to have declined from earlier periods (George Losey, pers. comm.).




Figure 19. Ta'ape abundance on Transects and 10 minute free swim surveys

Roi

Of the six species of groupers (family Serranidae) introduced to Hawai'i only roi, *Cephalopholis argus* has become established. There were more roi introduced (n=2385) than any other grouper and it was the only species introduced to the Island of Hawai'i (400 fish from Moorea in 1956). It now occurs on all the main Hawaiian Islands and in low numbers on some of the Northwest Hawaiian Islands.

As evidenced by transect and free-swim data (Figure 15) overall roi abundance at West Hawai'i sites was increasing since at least 1999 to 2004. West Hawai'i retrospective studies at Hōnaunau and Ke'ei indicate that roi populations only began to increase in the 1990's, three decades after their initial introduction. Randall notes in 1987 that "This fish (roi) has not become abundant. It has not developed a population approaching that of its native stock in the Society Islands."

Since 2004 however there has been a marked downturn in observed overall roi abundance both on West Hawai'i transect (56% decrease) and free swim surveys (55% decrease) (Figure 54). These declines occurred at 20 of 23 surveyed sites. This recent decline may be related in part to an unusual fish die off in West Hawai'i which first became apparent in May 2006. At that time seven dead roi were found washed up on the beach at `Anaeho`omalua, North Kona (Travis Hall, pers. Comm.). Several other species were also noted at this time including several goatfish (*Mulloidichthys* sp.), a surgeonfish (*Acanthurus dussumieri*) and a moray eel. Over the next five months there were numerous reports of dead and dying fishes, typically floating or struggling at the surface, along a wide stretch of the West Hawai'i coastline. In most instances the fish had distended swim bladders which prevented still live fish from returning to the bottom. Individuals of three species (*C. argus*, *C. sordidus* and *A. olivaceus*) were observed underwater live but having difficulty maintaining equilibrium. Roi were by far the most commonly involved species in the die off incidents but a number of other species also perished comprising a wide range of families, feeding types and depth ranges (Table 10). Similar undocumented reports of floating fish (typically roi) were also received from Maui, O'ahu and Moloka'i.

Ten specimens of nine species were collected and sent to the National Wildlife Health Center, U.S. Geological Survey in Honolulu for necropsy. Diagnostic Case Report findings typically indicated swim bladder distension, a variety of incidental lesions and, in two cases, atrophy of the liver. No gross or microscopic lesions were considered severe enough to cause death and the cause of death remains unknown (Thierry Work, pers. Comm.).

Table 4. List of fishes collected or reported in West Hawai'i die off

Family	Species	Common Name
Acanthuridae	<i>Acanthurus dussumieri</i>	Eyestripe surgeonfish
Acanthuridae	<i>Acanthurus olivaceus</i>	Orangeband surgeon
Acanthuridae	<i>Acanthurus triostegus</i>	Convict surgeonfish
Acanthuridae	<i>Ctenochaetus hawaiiensis</i>	Black surgeonfish
Acanthuridae	<i>Naso hexacanthus</i>	Sleek unicornfish
Acanthuridae	<i>Zebrasoma flavescens</i>	Yellow tang
Balistidae	<i>Melichthys niger</i>	Black durgon
Balistidae	<i>Rhinecanthus aculeatus</i>	Lagoon trigger
Balistidae	<i>Rhinecanthus rectangulus</i>	Reef triggerfish
Chaetodontidae	<i>Chaetodon auriga</i>	Threadfin butterflyfish
Chaetodontidae	<i>Forcipiger flavissimus</i>	Forcepsfish
Kuhliidae	<i>Kuhlia sandvicensis</i>	Hawaiian flagtail
Lutjanidae	<i>Lutjanus kasmira</i>	Ta'ape (Blueline snapper)
Mullidae	<i>Mulloidichthys</i> sp.	goatfish
Muraenidae	<i>Gymnothorax</i> sp.	Moray eel
Scaridae	<i>Chlorurus sordidus</i>	Bullethead parrotfish
Scaridae	<i>Scarus rubroviolaceus</i>	Redlip parrotfish
Serranidae	<i>Cephalopholis argus</i>	Roi (Peacock grouper)
Serranidae	<i>Epinephelus quernus</i>	Hawaiian grouper




Figure 20. *C. argus* density in West Hawai'i. Data based on two types of underwater visual surveys at 26 long-term monitoring stations spread over approximately 100 miles of coastline. Each site was surveyed 4-6 times a year

The following year in 2007 only a single fish was reported or found suffering similar conditions, that being the deep-sea swallower *Kali indica* (Fig 55).



Figure 21. *Kali indica*

Puffer die-off

Early in 2010 a die-off of large puffers, with external symptoms quite similar to the previous mortalities, began to occur on Maui and Hawai'i Island. Over the ensuing months low numbers of dead and dying puffers increased (Figure 22) and were progressively reported up the island chain as far as Kaua'i (Oct. 2010). The overall reported numbers of dead puffers decreased as fall approached. Greater than 95% of all reported mortalities were of the stripebelly puffer, *Arothron hispidus* with a few porcupine fish (*Diodon hystrix*), Hawaiian Whitespotted toby (*Canthigaster jactator*) and spotted puffer (*Arothron meleagris*) (Thierry Work, pers. comm.)

A network of concerned citizens and agency people were actively involved in this incident, filing reports of mortalities and shipping dead fish to Dr. Thierry Work, Wildlife Disease Specialist with the U.S. Geological Service (USGS) in Honolulu who performed both gross and microscopic examinations. All assays for viruses (including electron microscopy) have so far come up negative and all attempts to incriminate any other infectious agent as a cause have come to naught. The current hypothesis is that the puffers died from a natural toxin of some sort. A chemist at the Charleston SC NOAA lab has been working the problem for the last 2 years. A natural toxin has been isolated from the puffers and the lab is in the process of trying to determine its structure.




Figure 22. Number of dead puffers examined at USGS Honolulu Field Station

West Hawai'i monitoring data indicates a substantial decline has occurred in the spotted puffer (*A. meleagris*) with a precipitous drop in 2009/2010 (Figure 23). Other large puffer species were too infrequently counted on transects to determine changes in abundance. The decline in *A. meleagris* is somewhat perplexing in that this species did not constitute a substantial portion of the reported and examined mortalities. It is of interest to note that two separate dead puffers of this species were found face down underwater at Ke'ei (photo in Fig 23) and in a Wai o Pae tide pool (Jennifer Turner, pers. comm.). In a somewhat similar vein, West Hawai'i monitoring data indicates that the Hawaiian spotted toby (*C. jactator*) declined substantially over the past decade. It's unknown whether similar sorts of mortalities are responsible.




Figure 23. Pufferfish abundance in West Hawai'i

Roi impacts

As previously noted, although roi was introduced to augment declining populations of food and game fishes it has not been well received by most Hawai'i fishermen due to concerns about ciguatera and more recently about negative impacts to native fish populations. As with ta'ape, roi have been blamed for a multitude of problems on the reefs, including a decline in important aquarium fish such as the yellow tang *Zebrasoma flavescens*. Concern has also been expressed over putative impacts on food fishes and invertebrates

The marked decline in the numbers of West Hawai'i roi in recent years provides an unprecedented opportunity (i.e. a 'natural' experiment) to examine responses of the reef fish community to a >50% reduction in the roi population. This work is currently planned for the coming year. It is anticipated that if roi are having major impacts on the abundances of other species they prey upon there would be detectable and consistent temporal relationships between roi and prey species abundance. An examination of roi and two of the most abundant species in roi's prime habitat the yellow tang (*Zebrasoma flavescens*) and kole (*Ctenochaetus strigosus*) fails indication direct negative impact on either species. From 1999 to 2004 as roi populations were increasing, both kole and yellow tang populations were increasing. Subsequent to 2004 as roi populations decreased yellow tans similarly decreased whereas kole numbers were fairly stable

(Figure 58). This is not the pattern that would result if roi abundance was a major determinant of the abundance of these other two species.




Figure 24. Temporal trends of the numbers of kole, yellow tang and roi in FRAs. Young of Year (YOY) not included

Another complementary way of examining the extent and magnitude of potential roi impacts on West Hawai'i reef fish populations is to examine the relationship between roi abundance at each of the monitoring sites with the abundance of various species and functional groups at the sites. Figure 25 illustrates this approach for six different groups of fish; none of which show a significant negative relationship with roi abundance. In other words having more roi in an area does not result in having less; A). total fish ($p=0.58$), B). small prey fish ($p=0.86$), C). other piscivores ($p=0.24$), D). yellow tang YOY (0.16), E). Kole YOY ($p=0.79$) or F). all YOY ($p=0.86$).




Figure 25. Relationship between Roi and various West Hawai'i fish population parameters

Aquarium species

The aquarium collecting industry in Hawai'i and especially in West Hawai'i has long been a subject of controversy. In contrast to other areas in the State, the West Hawai'i aquarium fishery has undergone substantial and sustained expansion over the past 35 years (Figure 24). Approximately 79% of fish caught in the State and 68% of the total aquarium catch value presently comes from the Big Island (Table 5).

Prior to the last two years, almost all of the Big Island catch (97% avg. from 2001 – 2010) came from West Hawai'i. However, in recent years, there has been a dramatic increase in the catch of *Halocaridina rubra* (opae ula), an endemic anchialine shrimp both from West and East Hawai'i (Figure 26). This shrimp is widely used in 'microcosm' displays (mini ecosystem environments) which are widely available for sale online. Although at least one company (petshrimp.com) claims to sell only captive bred opae ula, catch data indicates substantial numbers of wild-caught shrimp are currently being taken.




Figure 26. Number of marine aquarium animals collected and number of commercial aquarium permits in West Hawai'i for fiscal years 1976-2011

Table 5. Changes in West Hawai'i aquarium fishery since implementation of the FRAs. Dollar value is adjusted for inflation

	FY 2000	FY 2012	Δ
No. Permits	48	54	13% ↑
Total Catch	279,606	388,344	39% ↑
Total Value	\$745,129	\$1,184,610	59% ↑
% of State Fish Catch	70%	79%	9% ↑
% of State Fish Value	67%	71%	4% ↑
% of State Total Catch	55%	69%	14% ↑
% of State Total Value	59%	68%	9% ↑

The West Hawai'i Regional Fishery Management Area, which spans the entire West Hawai'i coastline, was established in 1998 primarily in response to controversy surrounding the activities of aquarium collectors working the coastline. Overall, the marine aquarium fishery in the State of Hawai'i is one of the most economically valuable commercial inshore fisheries with FY 2012 reported landings of 439,358 specimens and




Figure 27. Opae ula catch in East and West Hawai'i Keoki Stender photo

a total value of \$2.05 million. Earlier studies suggested that the reported values may have been underestimated by a factor of approximately 2 to 5X (Cesar et al. 2002, Walsh et al. 2003). However, a recent analysis (FY 2010) comparing Hawai'i Island aquarium catch report data with dealer purchase data from the collectors found good correspondence in reported numbers. There was only a 3.5% difference between the number of animals reported caught and sold by aquarium collectors. More importantly dealer reports of purchases from Hawai'i collectors did not indicate underreporting by the collectors (Figure 27). Walsh et al. 2003 provides an historical overview of the commercial aquarium fishery in Hawai'i.




Figure 28. Comparison between Hawai'i Island aquarium collector report data and dealer purchases of aquarium animals from the collectors (FY 2010)

In 1999, DAR in conjunction with a citizen's advisory group, the West Hawai'i Fisheries Council (WHFC), established a network of 9 Fish Replenishment Areas (FRAs) where aquarium collecting was prohibited. Along with existing protected areas 35.2% of the coastline was off limits to collecting.

In order to investigate the effectiveness of the FRAs to replenish depleted fish stocks, a consortium of researchers established the West Hawai'i Aquarium Project (WHAP) in early 1999. Funding was secured for the early years of the project through the Hawai'i Coral Reef Initiative Research Program (HCRI-RP), a federal initiative under the aegis of the National Oceanic and Atmospheric Administration (NOAA). Subsequent funding has been provided by Coral Reef Monitoring Grants under NOAA's Coral Reef Conservation Program. The initial project researchers were Dr. Brian Tissot, Washington State University, Dr. William Walsh, DAR/DLNR and Dr. Leon Hallacher, University of Hawai'i-Hilo. They have been joined in recent years by Dr. Ivor Williams, National Marine Fisheries Service, Dr. Mark Hixon, Oregon State University and Dr. Helen Fox, World Wildlife Fund.

WHAP established 23 study sites (Figure 12, Table 3) along the West Hawai'i coastline in early 1999 at 9 FRA sites, 8 open sites (aquarium fish collection areas) and 6 previously established Marine Protected Areas (MPAs) to collect baseline data both prior to and after the closure of the FRAs. The MPAs are MLCDs and Fishery Management Areas (FMAs), which have been closed to aquarium collecting for at least 9 years and were presumed to have close to "natural" levels of aquarium fish abundances. They serve as a reference or 'control' to compare with the FRAs and open areas. Three additional sites have been added over the years.

The overall goals of WHAP were two-fold: 1) To evaluate the effectiveness of the FRA network by comparing targeted aquarium fishes in FRAs and open areas relative to adjacent control sites and, 2) To evaluate the impact of the FRA network on the aquarium fishery.

The general rationale for WHAP's goals was based on the premise that changes in FRAs and open areas can best be estimated by comparing them to other areas which have been protected for relatively long periods of time. These areas (MPAs) serve as control areas against which the FRAs are measured both before and after the closure of the FRAs. This rationale is derived from a well-known statistical procedure known as the BACI (Before-After-Control-Impact) procedure (Tissot et al, 2004) which is an especially appropriate and statistically powerful method for examining FRA effectiveness.

For this study FRA effectiveness (R) is measured statistically as the change in the difference between each FRA and the mean of all MPA sites during each survey (control vs. impact) from before (1991-2000) vs. after (2007-2009) FRA establishment. Details on study methodology and this procedure are covered in (Tissot et al, 2004, Division of Aquatic Resources 2004).

R measures the changes within the FRA as a percent of the baseline abundance relative to control sites. In the case of this study, R is a measure of the effectiveness or 'protective value' of the FRAs. That is, what effect is increased protection having on targeted fish?

Scientific studies on reef fishes are notoriously difficult due to the very high variability of fish abundance in both time and space. Even with a rigorous statistical design (such as BACI) and 11 years of study, it is difficult to statistically detect changes in abundances except for the most common species that exhibit relatively large changes.

Fish Replenishment Areas (FRAs)

Changes in density for the ten most collected aquarium fishes across all FRAs are shown in Table 12. Yellow Tang density increased markedly (and significantly) in the FRAs while seven of 10 decreased (Achilles Tang, Multiband Butterflyfish and Brown Surgeonfish decreased significantly). However these seven species represent <6% of the total West Hawai'i aquarium catch.

The FRAs were 'effective' (increases in FRAs relative to long term MPAs) for eight of the top 10 collected species with three being statistically significant. As with density there were significant decreases in effectiveness for the Multiband Butterflyfish and Brown Surgeonfish. Both of these species are not very heavily collected averaging <2000 individuals per year over the last 5 years and are fairly abundant on the reef. It's thus not clear why their numbers are declining in the FRAs. These two species exhibited overall declines in all three types of areas with the greatest decrease in the protected areas (FRAs and MPAs). For the Brown Surgeonfish this may be the result of a competitive interaction with Yellow Tang and/or Goldring Surgeonfish (aka kole). As their numbers have increased the Brown Surgeonfish's has decreased. Both Yellow Tang and Brown Surgeonfish are herbivore browsers with quite similar diets (Jones 1968). In a possibly similar relationship Barlow (1974) found the numbers of Brown

Surgeonfish and manini (*Acanthurus triostegus*) to be negatively correlated and this was attributed to the aggressive dominance of the Brown Surgeonfish.

Table 6. Overall FRA effectiveness for the top ten most aquarium collected fishes. 'Before' = Mean of 1999-2000; 'After' = Mean of 2007-2009. YOY not included

COMMON NAME	SCIENTIFIC NAME	MEAN DENSITY (No/100M ²)		OVERALL% CHANGE IN DENSITY	ρ	R	ρ
		Before	After				
Yellow Tang	<i>Zebrasoma flavescens</i>	12.73	20.8	+63%	0.01	+77%	<0.01
Goldring Surgeonfish	<i>Ctenochaetus strigosus</i>	28.38	32.01	+13%	0.23	+83%	0.39
Achilles Tang	<i>Acanthurus achilles</i>	0.26	0.05	-81%	0.01	+2%	0.09
Orangespine Unicornfish	<i>Naso lituratus</i>	0.81	0.59	-27%	0.10	+2%	0.37
Chevron Tang	<i>Ctenochaetus hawaiiensis</i>	0.18	0.16	-12%	0.77	+3%	0.41
Longnose and Forcepsfish	<i>Forcipiger spp.</i>	0.64	0.84	+32%	0.13	+4%	0.03
Multiband Butterflyfish	<i>Chaetodon multicinctus</i>	5.20	3.49	-33%	0.02	-5%	<0.01
Brown Surgeonfish	<i>Acanthurus nigrofuscus</i>	8.58	4.06	-53%	0.03	-26%	0.01
Orangeband Surgeonfish	<i>Acanthurus olivaceus</i>	0.13	0.10	-20%	0.63	+3%	0.45
Ornate Wrasse	<i>Halichoeres ornatissimus</i>	0.94	0.65	-31%	0.08	+2%	0.14

Bold = statistically significant at $p \leq 0.05$

With only a single exception all of the FRAs have proven to be effective (positive R value) in enhancing Yellow Tang stocks (Figure 61). Seven of the eight increases were statistically significant. The single FRA which was ineffective was Waiaka'ilio Bay in North Kohala. This FRA had very low Yellow Tang recruitment throughout the study period and additionally the area may have been impacted by a sedimentation event in October 2006 on nearby reefs.

An examination of multiple factors associated with effective FRAs (Tissot et al., 2004) found that habitat quality, FRA size (especially reef width) and density of adult fishes are associated with significant recovery of fish stocks. Of particular importance are areas of high finger coral (*Porites compressa*) cover which is critical habitat for juvenile Yellow Tang and other fishes (Walsh, 1987). Live coral cover at Waiaka'ilio declined 12% between 2003 and 2007 (Table 1).




Figure 29. Effectiveness of individual FRAs to replenish yellow tang, 1999-2009.
 *= Statistically significant at $p \leq 0.05$

The overall average changes in yellow tang abundance in the three management areas are shown in Figure 29. Yellow Tang exhibited a delayed increase in abundance in all areas following a strong recruitment year in 2002. Relatively low recruitment in 6 of the following years resulted in subsequent downward trends in all areas. A similar pattern was evident with the high recruitment year of 2009 and subsequent lower recruitment years. Even with relatively low recruitment in 6 of the past 7 years the number of adult yellow tang has increased by 63% in the FRAs since they were established (Table 6).

When yellow tang reach sexual maturity they leave the deeper coral rich reef areas where they settled (and where DAR transects are located) for shallower reef habitat (Claisse 2009). For females this occurs at approximately 4-5 years of age and for males at age 5-7. Thus in the absence of substantial input of Young-of-the-Year fish, (i.e. low recruitment) yellow tang populations will invariably decline over time due to the emigration of mature fish in addition to natural mortality. Obviously such declines will be exacerbated by aquarium collection activities which specifically target smaller-sized fishes.




Figure 30. Overall changes in yellow tang abundance (Mean ± SE) in FRAs, MPAs and Open areas, 1999-2012. Yellow bars indicate mean density (May - Nov) Yellow Tang Young-of-Year (YOY). YOY are not included in trend line data

The decrease of yellow tang in open areas to below baseline levels is attributable to the above factors as well as an increase in the number of aquarium collectors and collected animals relative to the period when the FRAs were established (Figure 25). The continuing decline of yellow tang in areas open to collecting has prompted several additional proposed management actions including some size/bag limits, restricting which species can be collected (See Species of Special Concern section pg. 54) and the proposed establishment of a limited entry program for the fishery. Recruitment in 2009 was the highest in the past 10 years which is likely to ameliorate the overall downward trend for yellow tang in open areas, at least over the short term.

The fishing/reserve (i.e. FRA/MPA) impacts described above are striking, but of greater significance to the role such reserves have in enhancing and sustaining West Hawai'i populations and the fishery which depends on those, are effects of the reserve network on Yellow Tang breeding stocks. Based on adult yellow tang 'jet boot' surveys (Williams et al. 2009) it was found that adult densities were highest within protected areas and in 'boundary' areas (open areas adjacent to protected areas). Densities were lowest in open areas far from protected areas. The high densities in boundary areas are evidence of 'spillover' (outward movement from reserves into surrounding open areas) and indicate that protected areas supplement adult stocks not only within their own boundaries, but also in open areas up to a kilometer or more away. Thus, the 35% of the coastline in reserves helps to sustain yellow tang breeding stocks in about 50% of the coastline. There were no significant differences (Figure 30 t-test $p=0.71$) in the

abundance of adult yellow tang in open vs. closed areas based on shallow water (10'-20' depths) jet boot surveys (2006-2010). Total estimated coastwise population of adult yellow tang in this depth range was estimated to be >2.5 million individuals.




Figure 31. Yellow tang abundance in West Hawai'i shallow water (10'-20') habitat

Goldring surgeonfish or kole exhibited trends (Figure 63) quite similar to yellow tang but since they are more abundant and much less collected than the tangs, open areas have been relatively stable.




Figure 32. Overall changes in goldring surgeonfish (aka kole) abundance (Mean ± SE) in FRAs, MPAs and Open areas, 1999-2012. Bars indicate mean density (June-Nov) of goldring surgeonfish Young-of-Year (YOY). YOY are not included in trend line data

Overall, kole have increased by 13% since FRA establishment (Table 3). As with yellow tang, recruitment levels have been relatively high thus enabling densities to increase in the protected areas. It is unknown at present if kole make a habitat change as they reach sexual maturity. Recruitment patterns are markedly similar between the two species, likely due to similarities in spawning seasonality, location and daily timing (Walsh 1984, 1987).

Achilles tang (Figure 32) has generally shown a highly variable pattern in all management areas in the early years of the study with an overall decline in the last seven years. Average densities of this species is very low ($\bar{x} = 0.22/100m^2$) on all transects. The deeper reef areas where the DAR transects are located is not the prime habitat for adults of this species. They prefer the high energy shallower surge zones more typical of the shoreline drop-offs areas in West Hawai'i. Presumably algal food resources are more abundant in these areas. These areas reef areas are surveyed by means of the shallow water resource surveys conducted by DAR. Initial results from this program and other ancillary longer terms studies suggest there should be concern for the sustained abundance of this species. Achilles Tangs are a very popular food fish as well as an aquarium fish and thus are being harvested both as juveniles and adults. Low levels of recruitment over the past 11 years (\bar{x} (Jun-Nov) = $0.09/100m^2$) appear insufficient to compensate for the existing levels of harvest.




Figure 33. Overall changes in Achilles tang abundance in FRAs, MPAs and Open areas, 1999-2012. Bars indicate mean density (June-Nov) of Achilles tang Young-of-Year (YOY). YOY are not included in trend line data

DAR is currently in the process of developing a comprehensive package of size and bag limits for a number of popularly targeted species. There is a proposed bag limit of 10 Achilles tang/person/day which would only apply to aquarium collectors. Analysis of

aquarium catch report data indicates that such a bag limit will be largely ineffective in stemming the continuing decline in Achilles Tang since it is presently difficult to collect more than 10 Achilles tang in a day.

The abundance/recruitment trends of the orangespine unicornfish and chevron tang, the fourth and fifth most collected species, are somewhat similar to Achilles tang (Figures 33 & 34). Here again the primary adult habitat is not the deeper, coral rich areas, where the DAR transects are located. Additionally the orangespine unicornfish is also widely taken as a food fish as well as being an important aquarium fish. The abundance of both these species on the transects closely tracks recruitment with an upturn during 2004/2005 when there was somewhat higher recruitment followed by declining trends in subsequent years that had low recruitment. Overall, recruitment has been minimal over the last 14 years for both orangespine unicornfish ($\bar{x} = 0.03/100m^2$) and chevron tang ($\bar{x} = 0.05/100m^2$).




Figure 34. Overall changes in orangespine unicornfish abundance (Mean \pm SE) in FRAs, MPAs and Open areas, 1999-2012. Bars indicate mean density (June-Nov) of clown tang Young-of-Year (YOY). YOY are not included in trend line data




Figure 35. Overall changes in chevron tang abundance (Mean ± SE) in FRAs, MPAs and Open areas, 1999-2012. Bars indicate mean density (June-Nov) of chevron tang Young-of-Year (YOY)

As observed in previous work (Walsh 1987) and emphasized again in this work, for some species, recruitment can be highly variable between years and repeated low levels of recruitment is a regular occurrence. Without substantial input of the YOY, overall abundances on the deeper reef transects decrease over time due to ontogenetic movement out of settlement habitat and natural mortality. This decrease can occur even in areas which are not subject to aquarium collecting pressure (i.e. FRAs and MPAs).

Although only a few species comprise the bulk of the West Hawai'i aquarium fishery, over 200 different species of fishes and invertebrates have been collected from the reefs over the last five years. Some of these species are uncommon or even rare and presumably have a low resilience to harvesting pressure. Even in protected areas a considerable amount of time may be required for populations of these species to increase. A good example seems to be the flame angelfish, *Centropyge loricula*. This very attractive but uncommon species is highly desired in the aquarium trade. Demand far exceeds the supply Hawai'i can provide so substantial numbers of this species are imported to Hawai'i (for subsequent reshipping) from other locales (e.g. Christmas Island).

Flame angelfish were rarely sighted on transect or free swim surveys during the first seven years of the study (Figures 35 & 36). Beginning in 2006 however they have become noticeably more abundant presumably due to one or more years of good recruitment although the recruits are apparently cryptic so not readily surveyed.

Flame angelfish abundance in open areas is decreasing in recent year presumably due to aquarium collection activities.




Figure 36. Flame angelfish on transects




Figure 37. Flame angelfish on free swim surveys

Species of Special Concern

Coral reef animals have multiple values and they serve fundamental biodiversity and ecosystem functions. They're important not only to aquarium collectors and other fishers but also to the commercial ocean recreation industry, their visitors and Hawai'i ocean users in general. Management of this resource needs to balance these values and uses. A number of reef fish species are particularly vulnerable to depletion because they may be naturally uncommon or rare but command high prices in the aquarium trade and are thus highly sought after by collectors. Examples include the dragon moray (*Enchelycore pardalis*), Tinker's butterflyfish (*Chaetodon tinkerī*), and bandit angelfish (*Apoemichthys arcuatus*). All of these species (and others) are worth more (sometimes considerably more) than \$50 each when collected. In a retail aquarium shop in Connecticut several years ago the author observed a bandit angel that sold for \$3,500.

For uncommon or rare species or those that occur in deeper reef habitats, it is difficult and/or unfeasibly expensive to gather solid information on their status and trends. Nevertheless for some of these species such as the Hawaiian turkeyfish there is considerable anecdotal evidence that they have declined in recent decades. It's also clear from a number of our long term studies at Puakō, Ke'ei and Hōnaunau that a number of fairly conspicuous species have likewise declined in abundance over time – most obviously several species of butterflyfish and, in particular, the bandit angelfish.

FRAs are a key component of the sustainable management of the West Hawai'i aquarium fishery. They encompass many of the areas most utilized by residents and dive/snorkel business, and help maintain the biodiversity of our reefs people expect and visitors are willing to pay for. The FRAs do not of course provide protection for species in the open areas. While they do provide a population reservoir, intensive fishing pressure on species with low natural abundances across most of West Hawai'i's reefs is problematic. Concerns over continued expansion of the fishery (up 25% in the last

decade) and harvesting effects in the open areas (65% of the coast), necessitate additional management measures.

To address such issues DAR in conjunction with The West Hawai'i Fisheries Council (WHFC) developed a 'white list' of species which could be taken by aquarium fishers (Table 7). The approach taken by the WHFC was based on the fact that the West Hawai'i aquarium fishery is very heavily focused on a relatively small number of species. Six species (yellow tang, goldring surgeonfish, Achilles tang, clown tang, black surgeonfish and Tinker's butterfly) make up 96% of the total catch value averaged over the last 5 years. The 40 species on the white list make up 99% of the total catch value so the great majority of species taken (over 180 species) have very little individual or collective value; nonetheless they are important components of the reef ecosystem. It should be noted no invertebrates are included on the white list.

The white list is part of a Hawai'i Administrative Rule (HAR 13-60.4) that is currently being processed. Although the list has been recommended and supported by the WHFC and approved by the nascent Big Island Association of Aquarium Fishers (BIAFF) there nevertheless has been some criticism directed at the list. Most of the concern is generally directed to why this species or that species is included on the list (i.e. allowed to be collected). Concerns have been articulated about collecting impacts on the species' populations and sometimes as to suitability and survivability of the species in captivity.

Aquarium Open vs. FRA Trend Analysis

In order to more comprehensively explore the 40 white list species and the current and potential impact to their populations on the reefs by aquarium collecting two different analyses were undertaken.

The first analysis examined the trends in the % difference in density between areas open to collecting and the FRAs (closed to collecting) for the species on the white list. Density was based on the overall average density of each species for the last three years (2010-2012) at all open and FRA survey sites. The % difference in fish densities between open and FRAs areas for a species was calculated as: $(\text{Density}_{\text{OPEN}} - \text{Density}_{\text{FRA}}) / \text{Density}_{\text{FRA}} \times 100$.

Table 7. Proposed 'White List' of species which can be taken by aquarium collectors within the West Hawai'i Regional Fisheries Management Area

Common Name	Scientific Name	Common Name	Scientific Name
Achilles Tang	<i>Acanthurus achilles</i>	Potter's Angelfish	<i>Centropyge potteri</i>
Goldrim Surgeonfish	<i>Acanthurus nigricans</i>	Pyramid Butterflyfish	<i>Hemitaurichthys polylepis</i>
Yellow Tang	<i>Zebrasoma flavescens</i>	Lei Triggerfish	<i>Sufflamen bursa</i>
Psychedelic Wrasse	<i>Anampses chrysocephalus</i>	Hi Dascyllus	<i>Dascyllus albisella</i>
Chevron Tang	<i>Ctenochaetus hawaiiensis</i>	Redbarred Hawkfish	<i>Cirrhitops fasciatus</i>
Milletseed Butterflyfish	<i>Chaetodon miliaris</i>	Hi Whitespotted Toby	<i>Canthigaster jactator</i>
Forcepsfish	<i>Forcipiger flavissimus</i>	Thompson's Surgeonfish	<i>Acanthurus thompsoni</i>
Fourspot Butterflyfish	<i>Chaetodon quadrimaculatus</i>	Saddle Wrasse	<i>Thalassoma duperrey</i>
Orangespine Unicornfish	<i>Naso lituratus</i>	Brown Surgeonfish	<i>Acanthurus nigrofuscus</i>
Yellowtail Coris	<i>Coris gaimard</i>	Black Durgon	<i>Melichthys niger</i>

Shortnose Wrasse	<i>Macropharyngodon geoffroy</i>	Fourline Wrasse	<i>Pseudocheilinus tetrataenia</i>
Gilded Triggerfish	<i>Xanthichthys auromarginatus</i>	Eightline Wrasse	<i>Pseudocheilinus octotaenia</i>
Goldring Surgeonfish	<i>Ctenochaetus strigosus</i>	Bluestripe Snapper	<i>Lutjanus kasmira</i>
Spotted Boxfish	<i>Ostracion meleagris</i>	Peacock Grouper	<i>Cephalopholis argus</i>
Orangeband Surgeonfish	<i>Acanthurus olivaceus</i>	Eyestripe Surgeonfish	<i>Acanthurus dussumieri</i>
Smalltail Wrasse	<i>Pseudojuloides cerasinus</i>	Tinker's Butterflyfish	<i>Chaetodon tinkeri</i>
Blackside Hawkfish	<i>Paracirrhites forsteri</i>	Blacklip Butterflyfish	<i>Chaetodon kleinii</i>
Bird Wrasse	<i>Gomphosus varius</i>	Fisher's Angelfish	<i>Centropyge fisheri</i>
Multiband Butterflyfish	<i>Chaetodon multicinctus</i>	Flame Wrasse	<i>Cirrhilabrus jordani</i>
Ornate Wrasse	<i>Halichoeres ornatissimus</i>	Hi Longfin Anthias	<i>Pseudanthias hawaiiensis</i>

There were 6 species which had distributions and/or behaviors which precluded obtaining accurate density estimates in the survey areas. *Chaetodon Kleinii* is a planktivore which typically feeds above the reef often near drop-offs or in deeper water. *Lutjanus kasmira* is a schooling species more likely to be found in deeper water at reef/sand interfaces while *Centropyge fisheri*, *Chaetodon tinkeri* and *Pseudanthias hawaiiensis* inhabit deeper (generally >50') waters. *Acanthurus dussumieri* were rarely recorded on fixed line transects and appeared to be associated with sand areas. Individuals of this species which are encountered are invariably of very large size and small fish (e.g. YOY) are rarely if ever seen on survey dives. These six species were excluded from the analyses.

The results of this analysis are presented in the following graphs (Figures 37-39). Given the controversial nature of all aspects of managing the aquarium fishery and the current relevance of the issue, available data for all 34 species are presented.

The columns (bars) represent the % difference in density between open and FRA areas for each year since 1999. Bars *below* the x axis indicate densities which are lower in the open areas relative to the FRAs and similarly bars *above* the x axis indicate densities which are higher in the open areas relative to the FRAs. The number to the right of the species name represents the 3 year (2010-2012) % difference. Note that 1999 data are prior to FRA establishment thus no FRA impact would be evident – rather just site differences between open areas.

The white list species can be classified into three groups based on their densities in the open areas relative to FRAs. Note that Young of Year are not included in these analyses. Group1 species (6 spp., Fig 37) had fairly consistent lower densities in the open areas. The yellow tang, *Zebrasoma flavescens* is particularly noteworthy as the disparity between the open areas and the FRAs is substantial. Averaged over the past three years (2010-2012) yellow tang are 61% less abundant in the open areas as compared to the FRAs. Yellow tangs are by far the most heavily targeted species in West Hawai'i and over the past decade the numbers of aquarium collectors and collected fish have increased substantially (Fig. 25). Although the disparity in yellow tang abundance was consistently increasing from 2000 to 2008 this trend has ameliorated somewhat over recent years.

The second most collected species, the kole, *Ctenochaetus strigosus*, also exhibits a collecting impact but in contrast to yellow tang the disparity between open and protected areas has not been increasing. For kole, open areas contain 32% fewer fish than the FRAs. For the multiband butterflyfish, *Chaetodon multicinctus*, the difference between

FRAs and the open areas has been consistently decreasing over the years and presently there are now slightly more of this species in the open areas than in the FRAs. Roi, *Cephalopholis argus* is, also less abundant in the open areas but this is not due to aquarium collecting as very few individuals of this species are collected (Table A). There is some indication that aquarium collectors kill this grouper on occasion or as a matter of course which may, in part, explain the difference between area types.

Group 2 species (11 spp. Fig., 38) did not exhibit any consistent pattern of differences in abundance in open vs. FRAs. In some years densities were higher in the FRAs while in other years they were higher in the open areas. In some years there were essentially no differences between areas. Aquarium collecting impacts, if any, are thus obscure.

Group 3 species (16 spp., Fig. 39) had consistently greater densities of fishes in the open areas vs. the FRAs. This pattern, as with Group 2 species, appears to relate to the comparatively low number of fishes collected relative to the size of their population on the reefs.

In summary, there was clear evidence of collecting impact for only 5 species of the 34 white list species which were analyzed. Four of the 5 (not *G. varius*) were all among the 10 most heavily collected species in the fishery (Walsh 2009). For the others, it appears that, at least based on the past 13 years data, inclusion on the white list poses little or no threat to their populations. The caveat is that this assumes collecting preferences will remain similar to the past decade and the amount of collecting effort (i.e. number of collectors) does not substantially increase. Furthermore these findings do not mean that aquarium collecting may not be having major impacts on species not on the white list especially uncommon, rare and valuable species.




Figure 38. White list species showing fairly consistent lower densities in areas open to aquarium collecting. The graph columns represent the % difference in density between open and FRA areas. Bars below the x axis indicate densities are lower in the open areas relative to the FRAs. 'E' denotes an Endemic species




Figure 39. White list species exhibiting inconsistent differences in density between areas open to aquarium collecting and FRAs. The graph columns represent the % difference in density between open and FRA areas. Bars below the x axis indicate densities are lower in the open areas relative to the FRAs. Note different Y axis scale for *M. niger* and *A. nigricans*




Figure 39 con't. White list species exhibiting inconsistent differences in density between areas open to aquarium collecting and FRAs. The graph columns represent the % difference in density between open and FRA areas. Bars below the x axis indicate densities are lower in the open areas relative to the FRAs. Bars above the x axis indicate densities are higher in the open areas relative to the FRAs. Note different Y axis scale for *P. cerasinus* and *M. geoffroy*




Figure 40. White list species exhibiting higher population densities in areas open to collecting relative to FRAs. The graph columns represent the % difference in density between open and FRA areas. Bars above the x axis indicate densities are higher in the open areas relative to the FRAs




Figure 40 con't. White list species exhibiting higher population densities in areas open to collecting relative to FRAs. The graph columns represent the % difference in density between open and FRA areas. Bars above the x axis indicate densities are higher in the open areas relative to the FRAs.




Figure 40 con't. White list species exhibiting higher population densities in areas open to collecting relative to FRAs. The graph columns represent the % difference in density between open and FRA areas. Bars above the x axis indicate densities are higher in the open areas relative to the FRAs.

Aquarium Species Population and Catch Analysis

The second approach to assessing white list inclusion estimated actual populations of the species on the list and related those numbers to the aquarium catch of that species. Most aquarium collecting in West Hawai'i occurs primarily in mid-depth ranges. While abundance and conditions can and will alter collecting depths, Stevenson et al. (2011) reported that the majority of aquarium fishers collect between 41' - 59'. A population estimate was thus made based on a depth range of 30'-60' which broadly corresponds to the depths encompassed by DAR West Hawai'i transect data (Table 3). An added advantage of this data set is that survey sites span a considerable portion of the West Hawai'i coastline and include both open and closed areas.

Mean densities for the 34 species on the white list for which adequate data existed were calculated for the period 2010-2012 at open survey sites. A GIS was used to determine the total area of hard bottom reef in the 30' -60' depth range that was open to aquarium collecting. Open areas at the extreme north and south parts of the West Hawai'i coast were excluded due to the remoteness of the areas and the difficulty of operating and collecting there. Total

populations in the 30' -60' depth range were the product of open area density X open area (7.08 km²). This population was then related to the average catch of the species for the period FY 2010-2012 (Table X).

Table 8. Population estimates and % of population taken by aquarium collectors of 'White List' species. 'E' indicates an endemic species, Catch is the average aquarium catch over FY 2010 - 2012 and Population is an estimate of total numbers of fish (excluding YOY) in collected open areas of hard bottom in 30'- 60' depths. Catch as % of Population is the % of the species' population in collected open areas taken annually by aquarium collectors

Scientific Name		Catch	30'-60' Population	Catch as % of Population
<i>Zebrasoma flavescens</i>		294,521	574,188	51.29%
<i>Ctenochaetus strigosus</i>	E	38,404	1,731,768	2.22%
<i>Acanthurus achilles</i>		9,801	8,496	115.36%
<i>Naso lituratus</i>		4,265	77,880	5.48%
<i>Ctenochaetus hawaiiensis</i>		2,602	13,452	19.34%
<i>Forcipiger flavissimus</i>		1,409	26,904	5.24%
<i>Chaetodon multicinctus</i>	E	1,292	233,640	0.55%
<i>Centropyge potteri</i>	E	1022	147,972	0.69%
<i>Halichoeres ornatissimus</i>		926	143,724	0.64%
<i>Acanthurus nigrofuscus</i>		809	940,224	0.09%
<i>Acanthurus olivaceus</i>		785	24,072	3.26%
<i>Chaetodon quadrimaculatus</i>		662	14,586	4.54%
<i>Thalassoma duperrey</i>	E	653	212,236	0.31%
<i>Coris gaimard</i>		614	9,833	6.24%
<i>Anampses chrysocephalus</i>	E	387	NA	NA
<i>Acanthurus nigricans</i>		381	3,278	11.62%
<i>Gomphosus varius</i>		338	37,694	0.90%
<i>Chaetodon miliaris</i>	E	313	7,375	4.24%
<i>Macropharyngodon geoffroy</i>	E	252	2,950	8.54%
<i>Pseudojuloides cerasinus</i>		244	14,094	1.73%
<i>Chaetodon tinker</i>		217	NA	NA
<i>Sufflamen bursa</i>		209	43,431	0.48%
<i>Hemitaurichthys polylepis</i>		181	27,861	0.65%
<i>Ostracion meleagris</i>		152	4,753	3.20%
<i>Dascyllus albisella</i>	E	149	37,203	0.40%
<i>Acanthurus thompsoni</i>		133	50,150	0.27%
<i>Pseudocheilinus octotaenia</i>		126	126,358	0.10%
<i>Canthigaster jactator</i>	E	97	143,075	0.07%
<i>Cirrhilabrus jordani</i>	E	96	NA	NA
<i>Pseudocheilinus tetrataenia</i>		81	207,319	0.04%
<i>Melichthys niger</i>		79	17,864	0.44%
<i>Pseudanthias hawaiiensis</i>	E	75	NA	NA
<i>Centropyge fisheri</i>		74	15,078	0.49%
<i>Acanthurus dussumieri</i>		61	NA	NA
<i>Chaetodon kleinii</i>		53	2,622	2.02%
<i>Lutjanus kasmira</i>		52	4,425	1.18%

<i>Paracirrhites forsteri</i>		45	11,964	0.38%
<i>Xanthichthys auromarginatus</i>		29	7,867	0.37%
<i>Cirrhitoys fasciatus</i>		9	5,081	0.18%
<i>Cephalopholis argus</i>		1	18,519	0.01%

Based on this analysis aquarium collecting is having the largest impacts on Achilles and yellow tang. For Achilles tang, the catch as a % of population in excess of 100% suggests that these tangs are being collected in areas apart from the open areas surveyed. Several collectors have indicated that fair numbers of Achilles tang still occur along the most southerly stretch of reefs on the Island of Hawai'i. Achilles tang has had low levels of recruitment over the past decade (Fig 32) and substantial numbers of larger fish (i.e. 'breeders') are taken for human consumption. Given these factors, population declines and a substantial aquarium impact are not surprising. There is currently a proposed bag limit for aquarium collectors of 10 fish/person/day undergoing Hawaii Administrative rulemaking.

Yellow tang has generally recruited reliably (Figure 29) and aquarium take has been decreasing in recent years from a previous period of continual and likely unsustainable increases (Fig 40). The price per fish paid by dealers to collectors has increased almost 1.8X since 2000 but has declined over the last three years, likely an effect of the U.S. economic recession.




Figure 41. West Hawai'i Yellow tang catch since FRA establishment and price per fish (adjusted for inflation)

For most of the species on the white list collecting impact, in terms of the % of the population being removed annually, is relatively low with 12 species having single digit % catch and 19

species having % catch values <1%. It should also be noted that the % catch does not include targeted fishes which occur in waters shallower than 30' and deeper than 60'. As such the estimates overestimate the actual % take of the population for many, if not most, species.

Endemic Species on the WHFC White List

An endemic species is a one whose presence is restricted to a defined geographic area. Of the 662 species of reef and shore fishes in the Hawaiian Islands it is currently estimated that 25% of them are endemic (Randall 2007). Many species endemic to the Hawaiian Islands also occur at Johnston Atoll. A number of Hawaiian endemics are important food fishes and are harvested both commercially and non-commercially. These include such fish as manini, āholehole, 'alai'ihi 'āweoweo, hāpu'u, kole, kūmū, mamo, nabeta, nohu, uhu and 'upāpalu and spiny lobsters and all opihi.

Several researchers have commented on the relative abundance of endemic fishes. Gosline and Brock (1960) noted "that many of the endemic fish of the Hawaiian Islands are the most abundant of their genera" and similarly Hourigan & Reese (1987) state that "many endemic species are the most abundant Hawaiian fishes in their families". Randall (2007) commented that "native species have evolved in isolated outposts such as Hawaii for long periods of time and therefore have had ample opportunity to become fully adapted to their environment".

Of the 40 species on the WHFC white List, 11 (27.5%) are considered endemic to Hawai'i - only slightly about the average level of overall endemism. All but one (*Anampses chrysocephalus*) also occurs at Johnston Atoll. The endemic white list species are listed in the table 9. Notes to relative abundance are referenced below. Listed in the third column are population estimates on West Hawai'i reefs in hard bottom habitat in 30'-60' depths. These estimates are derived from WHAP survey densities (2010-2012) and area estimates from NOAA habitat maps. The fourth column lists the % of a species population in 30'-60' Open areas which is taken annually by aquarium collectors (based on FY 2010- 2012 records).

Table 9. Endemic species on 'white list'

Species	Notes	30'-60' WHI Pop	% AQ Catch 30'-60' Open Areas
<i>Canthigaster jactator</i>	Most common Toby ¹	114,447	0.2%
<i>Thalassoma duperrey</i>	Most common inshore wrasse ¹	548,014	0.3%
<i>Centropyge potteri</i>	Most common angelfish ¹	310,666	0.6%
<i>Dascyllus albisella</i>		56,512	0.6%
<i>Chaetodon multicinctus</i>		661,184	0.6%
<i>Ctenochaetus strigosus</i>	Very common on HI reefs ¹	5,334,800	2.0%
<i>Chaetodon miliaris</i>	Most common B-Fly ^{1,2}	9,347	4.1%
<i>Macropharyngodon geoffroy</i>		2,864	6.5%
<i>Anampses chrysocephalus</i>		1,854	8.7%
<i>Cirrhilabrus jordani</i>	Common in right habitat ³	NA*	NA
<i>Pseudanthias hawaiiensis</i>	Abundant at 40-199m ⁴	NA	NA
NA* - Species occurs in habitats deeper than transects			

¹ Randall, JE. 2007, ² Brock, VE and TC Chamberlain. 1968, ³ Hoover, JP. 2008, ⁴ Chave, EH and BC Mundy. 1994

figures 37-39 presented the difference in a species' abundance in West Hawai'i Fish Replenishment Areas (FRAs, n=9) relative to Open Areas (n=9). Bars represent the % difference in abundance for each year from 1999 to 2012. Bars above the horizontal x axis

indicate the species was more abundant in the Open Areas (aquarium collected) than the FRAs. Similarly, bars below the x axis indicate greater abundance in the FRAs than the Open Areas.

Of the nine endemic species for which we have survey data, only two species were consistently less abundant in the Open Areas. Kole (*Ctenochaetus strigosus*) relative abundances have been fairly consistent over the past 14 years being approximately 30% less abundant in Open Areas than in FRAs (avg. 2010-2012). The multiband butterflyfish (*Chaetodon multicinctus*) has also been consistently less abundant in the Open Areas but this difference has been decreasing in recent years and now there are now slightly more of this species in the open areas than in the FRAs. (avg. 2010-2012). The % of the population of both these species taken by aquarium collectors in recent years is 2% or less (see Table).

Six of 11 endemic species on the white list are regarded as being common in suitable habitat. The population estimates presented represent only a portion of available habitat where these species occur. Thus actual populations are invariably higher than indicated for just the 30'-60' depth range.

For the 9 species for which we have data all of them have <9% of their open area population collected annually. Five of the 10 species have <1% of their population collected. Populations in MPAs and FRAs are essentially not collected and as indicated above total populations are higher than estimated in just the 30'-60' depth range. This means the percentage of total population taken by aquarium collectors is substantially lower than indicated in the table above. Only two of nine endemic species are consistently less abundant in Open Areas relative to FRAs suggesting very low aquarium related impact on the other species at present. While the difference in relative abundance has been stable for kole it has been decreasing for the multiband butterflyfish and the portion of the population taken by aquarium collectors is very low.

Given past and present collecting preferences and effort, the inclusion of these endemic species on the white list appears to pose little or no threat to populations on West Hawai'i reefs. Attention should continue to be paid to the psychedelic wrasse (*A. chrysocephalus*) and the shortnose wrasse (*M. geoffroy*) which have the lowest estimated populations in the depth range and the highest relative level of collection. Similarly caution should be exercised regarding those species for which abundance data is lacking.

Aquarium reef fish catch vs. non-aquarium catch

Controversy over aquarium collecting has become ever most pervasive in recent years due primarily to a small cadre of anti-aquarium collecting zealots on the island of Maui. In their view, management of the aquarium fishery is not an option; it should not even be regarded as a 'fishery. Only a total outright ban is acceptable. Unfortunately their concern regarding impacts of aquarium collecting has not focused solely on the reefs of Maui. Considerable time, effort and expense has been expended by this group at thwarting community-based management efforts in West Hawai'i such as the establishment of the 'white list, size and bag limits for key targeted aquarium species and a West Hawai'i specific aquarium permit. The latter is a preliminary step in the development of a limited entry aquarium fishery along this section of the island.

In order to gain a more balanced perspective on the generalized impact on reef fishes by aquarium collecting vis á vis other types of reef fishing activities, reef fish landings by aquarium collectors were compared with that of other commercial fishers and non-commercial 'recreational' fishers. Both aquarium collectors and other commercial fishers are required by

law to submit catch reports and thus island specific reef fish catch data is available for each group. As noted previously (Fig XX) recent analysis suggests that aquarium catch reports appear to fairly accurately reflect actual catch. Unfortunately similar assurance isn't available for other commercial catch reports.

Recreational fishers in Hawai'i are not required to submit catch reports but such catch data has been collected since 2003 by the Hawaii Marine Recreational Marine Fishing Survey (HMRFS) and subsequently in 2007 by NOAA's Marine Recreational Information Program (MRIP). Species-specific recreational catch data is available online (http://www.st.nmfs.noaa.gov/st1/recreational/queries/custom_time_series.html) on a statewide basis. All catch data from 2008 thru 2010 was decreased by a factor of 81.96% (i.e., 1/1.22) because of a count error in the population household numbers for Maui County (Hongguang 2012).

Over the past four years the number of reef fishes caught statewide by the recreational and commercial sectors been quite comparable averaging 1,511,025/yr. for recreational fishers and 1,554,010/yr. for commercial (i.e. non-aquarium) fishers Fig. 41). The




Figure 42. Comparison of the number of reef fishes caught by recreational and commercial fishers in the Main Hawaiian Islands

combined catch however is 1.7X the total statewide take (1,810,402/yr.) of aquarium fishes. The average yearly biomass (pounds) of reef fish caught by commercial fishers was similar for both commercial fishers (1,199,520 lbs.) and recreational fishers (1,160,337 lbs.) (Fig. 42). A biomass comparison was not made with the aquarium catch.




Figure 43. Comparison of the biomass of reef fishes caught by recreational and commercial fishers in the Main Hawaiian Islands

To compare total reef fish catches for the various fishing sectors on a more localized area basis it was necessary to apportion the recreational catch among areas. An adjustment factor was calculated based on the percentage of statewide commercial reef fish landings reported from each area (generally island or county as well as West Hawai'i). A separate adjustment factor was derived for both number of reef fishes caught and biomass. Biomass was estimated for aquarium fish catch by specifying a targeted size or typical maximum size of collected species based on information provided by active collectors (n= 7) and Stevenson et.al. (2011). Size data was then converted to weight utilizing length to weight conversion factors (DAR database).

In West Hawai'i the aquarium fishery takes 1.8X the number of reef fishes taken by recreational and other commercial fishers (Figure 44). 90% of the aquarium caught fishes are a single species – the yellow tang. In terms of all other reef fish species, the recreational and commercial fisheries combine to take 3X the number of reef fishes caught by aquarium collectors (Figure 45).




Figure 44. Comparison of the number of reef fishes caught by recreational, commercial and aquarium fishers in West Hawai'i




Figure 45. Comparison of the number of reef fishes, excluding yellow tang, caught by recreational, commercial and aquarium fishers in West Hawai'i

In terms of reef fish biomass caught by the different fisheries in West Hawai'i, considerably more biomass is taken by the combined recreational and commercial fisheries either including yellow tang (2.8X) or excluding it (8.6X) (Figures 46 & 47). Additionally, unlike the aquarium fishery which targets mostly immature fish, the other fisheries selectively target the larger breeding portion of the population which has profound implications for the sustainable usage of the resource.




Figure 46. Comparison of the biomass of reef fishes caught by recreational, commercial and aquarium fishers in West Hawai'i




Figure 47. Comparison of the biomass of reef fishes, excluding yellow tang, caught by recreational, commercial and aquarium fishers in West Hawai'i.

On Maui where, as noted, there has been considerable concern over putative aquarium collecting impacts the numbers of reef fishes caught by recreational and commercial fishers is 22X that taken by aquarium collectors (Figure 48). In terms the biomass the differential is 145X (Figure 49). The total take of non-aquarium reef fish currently is substantially greater on Maui than it is in West Hawai'i (Table 10)




Figure 48. Comparison of the numbers of reef fishes caught by recreational, commercial and aquarium fishers on Maui




Figure 49. Comparison of the biomass of reef fishes caught by recreational, commercial and aquarium fishers on Maui

Table 10. Comparison of the number and pounds of reef fishes caught by recreational and commercial fishers on Maui and in West Hawai'i

	Maui		West Hawai'i	
	Recreational	Commercial	Recreational	Commercial
Number Caught	218,474	71,730	146,176	48,498
Pounds Caught	342,769	122,268	153,193	55,468

Gill net management

As mandated by Act 306, SLH 1998, a laynet (i.e. gill net) management plan was developed over four years by the WHFC and DAR. The recommended plan became administrative rule in 2005. The rule provides for continued small-scale subsistence-level netting while effectively controlling large-scale commercial netting. Eight areas have been designated where the use of gill nets is prohibited. Along with existing no gill-netting areas, approximately 25% of the coastline now prohibits the use of such nets (Figure 13).



Figure 50. Locations of laynet prohibited areas in West Hawai'i and shallow water resource fish survey sites

Additional provisions of the rule were designed to encourage responsible net use and enhance enforcement. These include requirements such as net registration and numbered identification (floats and tags), maximum soak time of four hours and maximum net length of 125'. One area (Kaloko-Honokōhau FRA) was designated a Hawaiian cultural netting area where only locally constructed handmade nets of natural fibers may be used. The West Hawai'i laynet rules served as a model for the rest of the state and have generally been adopted elsewhere except for Maui which completely banned their use

Transects conducted in shallow water habitats most likely to be impacted by lay gill netters (Figure 49) indicate there is presently little difference in the biomass of targeted food fishes between areas open to netting and those prohibiting netting either beginning in 2005 or MPAs which have had longer (>10 years) prohibitions on laynetting (Figure Z).




Figure 51. Biomass of 'Resource' (i.e. food) fish on shallow water transects. Only fish > 15 cm TL are censused. 'Open' denotes surveys (n=99) in areas where lay gill netting is permitted. 'New Gill Net' are survey areas (=32) which were closed to gill netting in 2005 and 'Long Term Gill Net' are survey sites (n=11) within MPAs which have prohibited netting for >10 years

The reasons for the lack of differences between open and laynet protected areas may relate to one or more of several factors: (i) the newly protected areas haven't had sufficient time to work; (ii) the protected areas are not effectively enforced; (iii) the sites of many of the shallow water resource transects may be areas where netting is impractical (i.e. rocky shorelines, sharp reef drop-offs, etc.) and (iv) the overall level of laynet fishing is relatively low. This last factor is supported by the low number of lay gill nets registered in West Hawai'i (52 as of Dec. 2009) as compared to the other islands (e.g. 796 on O'ahu).

Invertebrates

Crown of thorns

While *Acanthaster planci* is native to Hawai'i and not an introduced species it nevertheless is of substantial concern to the general public due to its reputation as a 'coral killer' and the publicity generated by massive outbreaks on other Pacific islands. The last reported large-scale occurrence in Hawai'i of the crown-of-thorns starfish, was in August 1969 when approximately 20,000 starfish were observed off the south shore of Moloka'i. Since that time there have only been scattered reports of COTS aggregations and all have been of considerably lesser magnitude. As noted previously, COTS have been implicated in recent coral declines on Maui.

Data from free-swim surveys reflect the low absolute abundance on the West Hawai'i reefs but does indicate an overall increasing trend in COTS abundance over the last seven years (Figure 51).




Figure 52. Overall Crown-of-Thorns abundance on West Hawai'i transects and 10 minute free swim surveys

Crown-of-Thorns Starfish Aggregation Monitored on Kona Coast

On September 13th, 2012, aggregations of *Acanthaster planci*, commonly known as the predator Crown-of-Thorn Starfish (COTS), were discovered at West Hawaii Aquarium Project (WHAP) Site 7, near Ka'ūpūlehu on the Kona Coast. This site is located at 10-18 m depth range.

One week later, on September 20th, as part of a scheduled monitoring survey, COTS were counted on four permanent 25m x 4m transects and during a 10 minute Rapid Visual Transect (RVT) survey around the perimeter of the site. To further assess the extent of the outbreak, four divers spaced approximately 10m apart at depths from 6m to 20m swam north from the site and conducted a 5 minute swim counting all COTS within a 5m visual belt survey. Due to the unusually high COTS abundance on the transects and the 5 minute swim, the team returned to the site on September 26, 2012 to further assess the outbreak. Surveys methods described above are part of a developing rapid response protocol established by Eyes of the Reef Network in collaboration with DAR monitoring techniques.

COTS counts were repeated along the permanent transects and RVT survey to compare to the previous week. COTS coral feeding scars (areas devoid of live tissue) were measured within a 4 meter belt along the 25m transect, along with coral colony size and species. The team then repeated the 5 minute swim north counting all COTS. These surveys indicated that the COTS aggregation had migrated farther north so another 5 minute swim count was conducted.

Over a one week period, *Acanthaster planci* (aka COTS) density decreased from 30 to 8 COTS per 200m² along the permanent transects at WHAP Site 7. Similarly, the 10 minute Rapid Visual Transect (RVT) survey around the perimeter of the site showed a decrease in abundance from 71 to 31 COTS. On September 20th, 5 minute timed surveys north of the site showed COT density at 58 per 200 m². Surveys one week later showed 45 COTS per 200m². An additional 5 minute survey continuing north revealed 45 COTS per 200m², with an extended survey determining additional aggregations of COTS were present further north and at shallower depths (<6m). Of two known COTS predators, *Cassia cornuta* and *Charonia tritonis* (aka Horned Helmet & Triton's Trumpet) only a single *C. tritonis* was observed throughout the surveys.




Figure 53. *Acanthaster planci* (COTS) density and abundances recorded on surveys conducted near Ka'ūpūlehu, WHAP Site 7 in September 2012. Zero COTS were observed on November 9, 2011

The most affected coral genus was *Montipora*, with approximately 131 colonies affected (average 82% of colony area dead), compared to 90% of area on 7 *Pavona* colonies, 85% of area on 30 *Pocillopora* colonies and 44% of area on 81 *Porites* colonies. (These are rough estimates as the scars were only measured along one axis).




Figure 54. Number of coral colonies observed at WHAP Site 7 with COTS feeding 'scars' and approximate area of subsequent colony mortality

During regular monitoring, zero COTS are usually observed along the four permanent transects and the RVT survey maximum is typically maxed out at a single individual (surveys conducted by DAR 4-6 times annually, beginning in 2000). However, on October 4, 2011, a total of 33 and 8 COTS were observed during the RVT and along the WHAP survey lines. Surveys conducted prior to and following the September 2012 outbreak, on July 31 and November 11, 2012) showed no COTS present at the site.

In September 2012, survey data shows an unusually high aggregation event of the predator *Acanthaster planci* (Crown-of-Thorns Starfish, aka COTS) for West Hawai'i. While the predominant coral species at this site is *Porites lobata*, the most affected genera were *Montipora* and *Pocillopora*. This finding is consistent with preferential feeding behavior on other Indo-Pacific reefs (Kayal et al. 2012). COTS aggregated in clusters with animals even piled upon one another rather than spread out (Figure 55A). There were well over 200 animals counted within an area where no more than a single individual is typically observed (WHAP data).

As mentioned above, COTS most frequently preferred Montiporid and Pocilloporid colonies, consuming over 80% of most colonies. Benthic surveys in 2011 along the permanent transects showed percent coral cover at this site for *Montipora sp.* and *Pocillopora sp.* to be 1% and 0.3% cover respectively. However, a substantial number of Poritids were also preyed upon, which comprise approximate 26% of the 27.1% total coral cover at the site. Tissue loss resulting from biological interactions such as COTS predation also has the ability to further influence benthic community structure and coral

health by making substrate available for algal colonization. Old COTS feeding scars were quickly overgrown by algae along the permanent transects (Figure 55B).

Tissue loss of corals is known to result from algal-coral interactions (Haas et al. 2010). Coral-algal interactions were surveyed along West Hawaii in 2010 and 2011 as part of a coral health monitoring program conducted by Courtney S. Couch (Cornell University Ph.D candidate) in collaboration with DAR. Algal overgrowth and the resulting coral tissue mortality were widespread both in the shallow habitats (3-6m) and deep habitats (WHAP sites). These surveys revealed that algal overgrowth was significantly higher in shallow habitats, with no clear seasonal trend across all the sites. Between 1-15% of all the coral colonies were overgrown by algae to some degree. Surveys revealed active algal overgrowth of live corals paired with tissue loss, primarily by *Corallophila huysmansii* (Figure 54C) (Couch et al. in prep).

C. huysmansii has the ability to settle on, overgrow, and kill live coral tissue through the hypothesized use of cytotoxic allelochemicals (Jompa & McCook 2003). With upwards of 15% of all colonies affected by algal overgrowth and approximately 44% of the affected Poritid colonies directly impacted by COTS predation at the outbreak site, biological interactions such as algal overgrowth and COTS predation may have a large, but underestimated, influence on not only coral health but also to benthic community structure and percent live coral cover along West Hawai'i.

Over several weeks of monitoring, the Crown-of-Thorns Starfish (COTS) aggregation appeared to be migrating slowly in a northerly direction and into shallower depths, presumably where more *Pocillopora* and *Montipora* colonies might be found, behavior which is consistent with other studies (Kayal et al. 2012). This COTS outbreak is clearly disturbing the coral community's diversity in an area that has already experienced strong storm damage and decreasing coral cover (net loss of 13% from 2003 - 2011). Moreover, only a single predator of COTS, *Charonia tritonis* (Triton's Trumpet), was observed at the site.

Monitoring of COTS populations will continue and immediate protection of their predators, *C. tritonis* and *Cassis cornuta* (Triton's Trumpet and Horned Helmet) is highly recommended throughout the State of Hawai'i.




Figure 55. A) Crown-of-Thorns Starfish (COTS) aggregation monitored at Ka'ūpūlehu (WHAP Site 7) in September 2012. Note: Arrow indicates COTS feeding scar on *Porites lobata* with algal colonization, B) Recent COTS feeding scar on *P. lobata* colony, C) Previously documented tissue loss of *P. lobata* colony at same site with colonization of turf algae and the filamentous *Corallophila huysmansii*, photographed on March 27, 2011

Urchins

Three of four of the most common surveyed urchin species have increased in West Hawai'i since monitoring began in 1999. Based on data presented in previous monitoring reports this increase does not appear to be related to a substantial increase in food supply (i.e. benthic algae) along the coast. Likewise there is no indication that potential food competitors such as herbivorous fishes (e.g. acanthurids) have markedly decreased. In actuality some of the most abundant surgeonfish have increased along with the urchins (Figure 56).

Even though urchin densities are increasing for three species, present abundance may still be lower than in previous years as can be seen at Kealakekua Bay which was surveyed for urchins in 1968 (Ebert, 1971).




Figure 56. Abundance (Mean \pm SE) of collector urchin *Tripneustes gratilla*, red pencil urchin *Heterocentrotus mammillatus* and banded urchin *Echinothrix calamaris* on transects

East Hawai'i

To date, abundance of fishes among sites is significantly different, being more abundant at both Waiopae sites than at Richardson's Ocean Center (Figure 57A). Species richness among sites is also significantly different among sites, being higher on MLCD transects compared to ROC (Figure 57 B). There are no among-site differences in species diversity ($p=0.435$) (Figure 57 C). The MLCD and ROC sites have the highest similarity in their fish communities, and the OPEN and ROC communities have the lowest similarity (Table 11).

(A)




Figure 57. Fish community parameters at Waiopae (MLCD & OPEN) and Richardson's Ocean Center (all survey years pooled) Data are means and standard errors. (A) abundance; (B) Species richness; (C) S-W Diversity.

Table 11. Percent Similarity from pairwise site comparisons.

Location	Percent Similarity
MLCD vs. OPEN	69.3%
MLCD vs. ROC	72.7%
OPEN vs. ROC	53.1%

Over the twelve years of surveying of fishes at Waiopae and Richardson's, there appears to have been a slight increase in fishes observed between 1999 and 2006, followed by a three-year decline, with an upturn on fishes seen so far in 2010 (Figure 57). There is generally good concordance in the year-to-year abundance of fishes among survey sites (Figure 57). Since the delineation of the Waiopae MLCD on June 16, 2003, no net increase in fish abundance has been observed.




Figure 58. Annual mean abundance (+SE) of fishes at Waioopae and Richardson's Ocean Center.

Of the 136 species recorded on transects at the three locations, most individuals are from one of six families: Labridae, Scaridae, Acanthuridae, Pomacentridae, Tetraodontidae, and Chaetodontidae (Table 11, Figure 58). Labrids and pomacentrids were particularly abundant at all three sampling areas, but scarids were only abundant on Waioopae Open transects. All of the transect lines in this area are deeper than other sites and two traverse a level area with abundant turf algae which appears to attract large numbers of scarids. Species densities at the three East Hawai'i sites are listed in Table X in Appendix.

Table 12. Individuals per 100 m² by family at East Hawai'i sites (n = 224 transects at Waioape Sites; n = 172 at Richardson's Ocean Center).

Family	OPEN	MLCD	ROC
Acanthuridae	13.10	6.12	9.88
Apogonidae	0.02	0.00	0.00
Aulostomidae	0.05	0.02	0.01
Balistidae	0.04	0.05	0.13
Belonidae	0.00	0.10	0.09
Blenniidae	1.30	1.06	0.35
Caracanthidae	0.00	0.01	0.04
Chaetodontidae	2.33	2.99	1.26
Cirrhitidae	0.11	0.40	1.41
Diodontidae	0.00	0.00	0.00
Fistulariidae	0.33	0.09	0.06
Gobiidae	0.03	0.01	0.00
Holocentridae	0.03	0.04	0.01
Kyphosidae	0.00	0.59	0.00
Labridae	48.54	52.46	39.52
Lutjanidae	0.04	0.40	0.00
Monacanthidae	0.02	0.01	0.01
Mugilidae	0.00	0.01	0.11
Mullidae	0.62	0.14	0.05
Muraenidae	0.09	0.14	0.14
Myliobatidae	0.00	0.00	0.00
Ophichthidae	0.00	0.00	0.00
Ostraciidae	0.05	0.19	0.05
Pomacanthidae	0.00	0.00	0.00
Pomacentridae	16.91	33.54	33.31
Scaridae	29.29	4.12	1.31
Scorpaenidae	0.00	0.07	0.23
Serranidae	0.15	0.41	0.01
Synodontidae	0.03	0.03	0.03
Tetraodontidae	4.12	4.11	2.72
Zanclidae	0.06	0.10	0.03
Pooled Individuals/100 m ² =	117.3	107.2	90.7




Figure 59. Wai Opae Open/MLCD and ROC fish abundance by family (all years pooled).

REFERENCES

- Aeby, G.S. (1998). A digenean metacercaria from the reef coral, *Porites compressa*, experimentally identified as *Podocotyloides stenometra*. *Journal of Parasitology* 84:
- Aeby, G.S. (2007). Spatial and temporal patterns of infection of *Porites* trematodiasis on the reefs of Kaneohe Bay, Oahu, Hawaii. *Bulletin of Marine Science* 80:209-218.
- Aeby, G.S, and Cotton, S.P. (2007) DAR monitoring report.
- Aeby, G., Williams, G.J., Franklin, E.C., Haapkyla, J., Harvell, C.D., Neale, S., Page, C.A., Raymundo, L.J., Vargas-Angel, B., Willis, B.L., Work, T.M. and Davy, S.K. (2011). Growth anomalies on the coral genera *Acropora* and *Porites* are strongly associated with host density and human population size across the Indo-Pacific. *PLoS ONE* 6(2): e16887. doi:10.1371/journal.pone.0016887.
- Baker, D. M. and Webster, K.L. (2010). Caribbean octocorals record changing carbon and nitrogen sources from 1862 to 2005. *Global Change Biology* 16(10): 2701-2710.
- Barlow, G.W. (1974). Extraspecific imposition of social grouping among surgeonfishes (Pisces: Acanthuridae). *J. Zool. Lond.* 174:333-340.
- Brock, V,E and T,C. Chamberlain. (1968). A geological and ecological reconnaissance off Western Oahu, Hawaii principally by means of the research submarine "Asherah". *Pac. Sci.* XXII: 373-394.
- Cesar, H., Beukering, P.V., Pintz, S., and Dierking, J. (2002). Economic valuation of the coral reefs of Hawai'i. Final Report. Hawai'i Coral Reef Initiative Research Program. 117 pp.
- Chave, E.H and B.C Mundy. (1994). Deep-sea benthic fish of the Hawaiian Archipelago, Cross Seamount and Johnston Atoll. *Pac. Sci.* 48(4): 367-406.
- Coral Disease Working Group. (2007) Coral disease, environmental drivers, and the balance between coral and microbial associates. *Oceanography* 20(1):172-195.
- Couch, C., S., Garriques, J., Barnett, C., Preskitt, L., Work, T.W., Giddens, J., Walsh, W. and Harvell, C. D. (in prep). Spatiotemporal Dynamics of Coral Health and Disease along the leeward coast of the Hawai'i Island. *Ecological Monographs*.
- Division of Fish & Game. Annual Report of the Board of Agriculture and Forestry, Territory of Hawaii. FY July 1,1955 – June 30,1956.
- Forsman, Z.H., Rinkevich, B., and Hunter, C.L. (2006). Investigating fragment size for culturing reef-building corals (*Porites lobata* and *P. compressa*) in *ex situ* nurseries. *Aquaculture*. 261:89-97.
- Grigg, R.W. and Maragos, J.E. (1974). Recolonization of hermatypic corals on submerged lava flows in Hawaii. *Ecology* 55: 387-395.

- Haas, A., el-Zibdah M., Wild C. (2009) Seasonal monitoring of coral-algae interactions in fringing reefs of the Gulf of Aqaba, Northern Red Sea.
- Harvell, C.D., Jordán-Dahlgren, E., Merkel, S., Rosenberg, E., Raymundo, L., Smith, G.W., Weil, E. and Willis, B.L. (2007). Coral disease, environmental drivers and the balance between coral and microbial associates. *Oceanography* 20:58-81.
- Hawaii Statewide GIS Program. Department of Business, Economic Development and Tourism Office of Planning. <http://hawaii.gov/dbedt/gis/>
- Hernandez-Munoz, D., Alcolado, P. M. and Hernandez-Gonzales, M. (2008). Effects of a submarine discharge of urban waste on octocoral (Octocorallia: Alcyonacea) communities in Cuba. *Revista de Biología Tropical*. 56(1):65-75.
- Hongguang, M. (2012). Catch and Effort Estimates for 2003-2010 from the Hawaii Marine Recreational Fishing Survey. PIFSC Internal Report IR-12-010. 08 March, 2012.
- Hoover, J.P. (2008). *The Ultimate Guide to Hawaiian Reef Fishes, Sea Turtles, Dolphins, whales and seals*. Mutual Publishing. 388p.
- Jompa, J., L.J. McCook. (2003) Coral-algal competition: macroalgae with different properties have different effects on corals. *Marine Ecology Progress Series*, Vol. 258: 87-95.
- Jones, R.S. (1968). Ecological relationships in Hawaiian and Johnston Island Acanthuridae (Surgeonfishes). *Micronesica* 4:309-361.
- Johnson, A. (2008) Groundwater discharge from the leeward half of the big island of Hawai'i. Masters in Science Thesis, University of Hawai'i.
- Kaczmarek, L. (2009) Characterizations of the major coral diseases of the Philippines: ulcerative white spot disease and novel growth anomalies of *Porites*. Biology, Miami.
- Kayal M., Vercelloni J., Lison de Loma T., Bosserelle P., Chancerelle Y., Geoffroy S., Stievenart C., Michonneau F., Penin L., Planes S., Adjeroud M. (2012) Predator Crown-of-Thorns Starfish (*Acanthaster planci*) outbreak, mass mortality of corals, and cascading effects on reef fish and benthic communities. *PLoS ONE* 7(10): e47363. doi:10.1371/journal.pone.0047363
- Knee, K.L., Street, J.H., Grossman, E.E., Boehm, A.B. and Paytan, A. (2010). Nutrient inputs to the coastal ocean from submarine groundwater discharge in a groundwater-dominated system: Relation to land use (Kona coast, Hawaii, USA). *Limnology and Oceanography* 55:1105-1122.
- Lough, J.M. and Barnes, D.J. (2000) Environmental controls on growth of the massive coral *Porites*. *Journal of Experimental Marine Biology and Ecology*. 245:225–243.
- Randall, J. E. (1987). Introductions of marine fishes to the Hawaiian Islands. *Bulletin of Marine Science*. 41:490-502.

- Randall, J.E. (2007). Reef and Shore Fishes of the Hawaiian Islands. Sea Grant Program. 546p.
- Richmond, R.H. (1987) Energetic relationships and biogeographical differences among fecundity, growth and reproduction in the reef coral *Pocillopora damicornis*. *Bulletin of Marine Science* 41(2):594-604.
- Rodgers, K., Cox, E. and Newton, C. (2003) Effects of mechanical fracturing and experimental trampling on Hawaiian corals. *Environmental management* 31(3):377-384.
- Tissot, B.N., Walsh, W. and Hallacher, L.E. (2004). Evaluating the effectiveness of a marine reserve network in Hawai'i to increase the productivity of an aquarium fishery. *Pacific Science* 58(2):175-188.
- Stevenson, T.C., Tissot, B.N. and Dierking, J. (2011). Fisher behaviour influences catch productivity and selectivity in West Hawaii's aquarium fishery. *Journal of Marine Science*. 68:813-822.
- Walsh, W.J. (1984). Aspects of nocturnal shelter, habitat space, and juvenile recruitment in Hawaiian coral reef fishes. Ph.D. Dissertation, University of Hawai'i. 475 pp.
- Walsh, W. J. (1987). Patterns of recruitment and spawning in Hawaiian reef fishes. *Environmental Biology of Fishes* 18(4):257-276.
- Walsh, W.J., Cotton, S.P., Dierking, J. and Williams, I.D. (2003). The Commercial Marine Aquarium Fishery in Hawai'i 1976-2003. In: Friedlander A.M. (ed.) *Status of Hawai'i's Coastal Fisheries in the New Millennium*. Proceedings of a Symposium sponsored by the American Fisheries Society, Hawai'i Chapter. pp. 132-159.
- Williams, G.J., Aeby, G.S., Cowie, R.O.M. and Davy, S.K. (2010). Predictive modeling of coral disease distribution within a reef system. *PLoS ONE* 5(2): e9264. doi:10.1371/journal.pone.0009264.
- Williams, I.D., Walsh, W.J., Claisse, J.T., Tissot, B.N. and Stamoulis, K.A. (2009). Impacts of a Hawaiian marine protected area network on the abundance and fishery sustainability of the yellow tang, *Zebrasoma flavescens*. *Biol. Cons.* 142:1066-1073.
- Yamashiro, H., Yamamoto, M and van Woesik, R. (2000) Tumor formation on the coral *Montipora informis*. *Diseases of Aquatic Organisms* 41:211-217.

Appendix A. Occurrences of eight coral diseases documented across 30 monitoring sites in West Hawai'i (GA = growth anomaly, TRE = trematodiasis, TLS = tissue loss syndrome, MFTL = multifocal tissue loss, HYP = hypermycosis).

ID	Site	<i>Porites</i> GA	<i>Porites</i> TRE	<i>Porites</i> TLS	<i>Porites</i> MFTL	<i>Pavona</i> HYP	<i>Montipora</i> GA	<i>Pocillopora</i> senescence reaction	<i>Pocillopora</i> TLS
SITE1	Lapakahi	x	x						
SITE2	Kamilo Gulch	x		x					
SITE3	Waiaka'ilio Bay	x		x					
SITE4	Puakō	x	x	x		x			
SITE5	Mauna Lani	x	x	x					
SITE6	Keawaiki		x			x			
SITE7	Ka'upulehu	x	x						
SITE8	Makalawena	x	x	x		x	x		
SITE97	Unualoha Pt.		x				x	x	
SITE9	Wawaloli Beach	x	x	x					
SITE10	Wawaloli FMA	x	x	x			x		
SITE11	Kaloko-Honokōhau	x	x	x					
SITE13	Papawai	x	x	x					
SITE98	Old Kona Airport	x	x	x					
SITE14	S. Oneo Bay	x	x	x		x			
SITE15	Keauhou	x	x	x					
SITE15x	Keauhou X	x	x	x		x			
SITE15y	Keauhou Y	x	x	x	x				
SITE15z	Keauhou Z	x	x	x					
SITE16	Kualani Pt. (Red Hill)	x	x	x	x		x		
SITE17	Red Hill	x	x						
SITE18	Keopuka	x		x					
SITE19	Kealakekua	x	x	x		x			
SITE20	Ke'ei	x	x	x	x				
HO	Hōnaunau drop off	x	x	x					
SITE21	Ho'okena (Kalahiki)	x	x	x	x				
SITE22	Ho'okena (Auau)	x		x				x	x
SITE23	Omaka'a Bay	x	x	x			x		
SITE99	Okoe Bay	x	x	x					
SITE24	Manukā	x	x	x					

**Appendix B. West Hawai'i Benthic Cover 2011 Surveys
Broad Benthic Categories**

Survey Site	Coral	Turf-Bare	Crustose Coralline	NCC Macroalgae	Macroalgae	Sand	Sessile Invert	Other
Lapakahi	11.8%	57.9%	0.5%	0.1%	0.1%	28.3%	0.0%	1.1%
Kamilo	29.0%	62.2%	5.8%	0.4%	0.1%	0.8%	0.0%	1.7%
Waiaka'ilio	38.8%	53.0%	4.8%	0.7%	0.8%	0.8%	0.0%	1.1%
Puako	34.2%	52.9%	9.6%	0.6%	0.4%	0.5%	0.0%	2.4%
'Anaeho'omalu	28.4%	57.1%	6.9%	0.9%	0.5%	2.4%	0.0%	3.8%
Keawaiki	18.7%	72.3%	5.1%	1.0%	0.1%	0.9%	0.1%	1.0%
Ka'upulehu	27.1%	62.3%	5.2%	0.6%	0.3%	2.8%	0.0%	1.7%
Makalawena	47.6%	49.3%	0.8%	0.0%	0.1%	1.3%	0.2%	0.9%
Unualoha	36.5%	59.4%	4.4%	0.2%	0.2%	0.7%	10.0%	1.7%
Wawaloli Beach	44.5%	52.0%	0.1%	0.0%	0.1%	1.3%	1.0%	1.0%
Wawaloli	42.3%	52.3%	0.2%	0.1%	0.1%	0.1%	3.6%	1.3%
Honokohau	48.3%	32.0%	1.9%	0.1%	0.0%	2.6%	13.0%	2.1%
Papawai	41.1%	46.9%	2.2%	0.5%	0.2%	0.9%	7.0%	1.2%
Old Kona Airport	51.2%	31.6%	1.2%	0.0%	0.0%	7.3%	7.6%	1.1%
S. Oneo	46.6%	43.9%	6.4%	0.2%	0.0%	1.1%	0.1%	1.8%
N. Keauhou	28.0%	64.5%	3.3%	1.5%	0.2%	0.1%	0.1%	2.5%
Kualani	62.4%	34.5%	1.3%	0.0%	0.1%	0.5%	0.0%	1.2%
Red Hill	35.3%	53.4%	3.2%	1.9%	0.9%	3.3%	0.1%	1.9%
Keopuka	14.4%	79.9%	3.2%	0.4%	0.2%	0.8%	0.0%	1.1%
Kealakekua	23.1%	64.8%	3.5%	0.1%	5.9%	0.0%	0.0%	2.6%
Ke'ei	26.7%	60.8%	3.1%	0.3%	6.3%	0.3%	0.0%	1.5%
Kalahiki	38.9%	45.4%	9.4%	2.2%	0.2%	1.9%	0.0%	1.2%
Au Au Crater	30.0%	56.0%	6.7%	3.5%	1.4%	0.2%	0.0%	2.2%
Omaka'a	32.9%	53.1%	5.7%	0.3%	0.8%	6.3%	0.0%	0.9%
Manuka	33.4%	52.7%	7.3%	2.7%	0.6%	1.7%	0.1%	1.5%

Appendix C. West Hawai'i Coral Cover By Species 2011 Surveys

Survey Site	<i>Montipora capitata</i>	<i>Montipora patula</i>	<i>Pavona varians</i>	<i>Pocillopora meandrina</i>	<i>Porites compressa</i>	<i>Porites evermanni</i>	<i>Porites lobata</i>	Other
Lapakahi	0.30%	0.00%	0.00%	0.30%	3.10%	0.60%	7.60%	0.10%
Kamilo	0.40%	0.30%	0.40%	0.10%	11.70%	0.00%	16.10%	0.00%
Waiaka'ilio	0.70%	0.20%	0.30%	0.30%	12.80%	0.00%	24.50%	0.00%
Puakō	0.20%	0.20%	0.20%	0.50%	12.00%	0.10%	21.20%	0.00%
Anaeho'omalu	1.10%	0.30%	0.40%	0.50%	9.20%	0.20%	16.60%	0.00%
Keawaiki	0.70%	1.40%	2.20%	0.30%	9.00%	0.00%	4.20%	0.90%
Ka'upulehu	0.60%	0.30%	0.00%	0.30%	2.20%	0.00%	23.80%	0.00%
Makalawena	1.50%	2.90%	1.60%	1.30%	8.70%	0.40%	24.40%	6.80%
Unualoha	0.50%	0.00%	0.00%	2.90%	3.40%	0.60%	28.50%	0.60%
Wawaloli Beach	0.8%	0.0%	0.0%	0.9%	6.1%	0.9%	35.7%	0.0%
Wawaloli	0.6%	0.0%	0.1%	3.7%	5.6%	0.2%	31.8%	0.4%
Honokohau	0.20%	0.00%	0.00%	0.10%	16.50%	0.90%	30.70%	0.00%
Papawai	0.50%	0.10%	0.00%	1.20%	3.50%	1.30%	34.20%	0.30%
Old Kona Apt	0.00%	0.00%	0.00%	0.70%	14.50%	0.80%	35.20%	0.00%
S. Oneo	0.10%	0.60%	0.60%	0.30%	17.40%	1.50%	26.10%	0.00%
N. Keauhou	0.10%	0.00%	0.00%	0.00%	16.80%	0.30%	10.80%	0.00%
Kualani	0.70%	0.20%	0.10%	0.40%	3.30%	13.90%	43.90%	0.00%
Red Hill	1.90%	0.30%	0.20%	0.80%	10.20%	0.50%	21.30%	0.00%
Keopuka	0.20%	0.20%	0.10%	2.20%	2.10%	0.30%	9.30%	0.00%
Kealakekua	0.10%	0.10%	0.40%	0.20%	7.00%	0.00%	15.40%	0.10%
Ke'ei	0.10%	0.00%	0.10%	0.10%	15.80%	2.90%	6.30%	1.50%
Kalahiki	0.20%	0.10%	0.00%	0.10%	16.40%	1.00%	21.30%	0.00%
Au Au Crater	2.40%	0.20%	0.10%	1.30%	5.60%	0.30%	20.00%	0.20%
Omaka'a	1.20%	0.10%	0.10%	1.10%	11.60%	2.10%	16.90%	0.00%
Manuka	0.20%	0.00%	0.10%	0.50%	8.90%	0.30%	23.30%	0.10%

Appendix D. West Hawai'i Benthic Cover 2007 Surveys

Broad Benthic Categories

Survey Site	Coral	Turf-Bare	Crustose Coralline	Encrusting Macroalgae	Macroalgae	Sand	Sessile Invert	Other
'Anaeho'omalu	31.5%	56.9%	7.1%	0.1%	0.3%	2.4%	0.1%	1.6%
Ho'okena	28.4%	57.7%	4.8%	2.3%	6.3%	0.1%	0.1%	0.3%
Honokōhau	48.5%	31.6%	3.3%	0.5%	0.3%	2.4%	12.9%	0.1%
Kalahiki	39.6%	48.3%	5.2%	2.2%	1.5%	2.0%	0.0%	1.2%
Kamilo	38.2%	51.0%	7.1%	0.3%	0.2%	0.7%	0.0%	2.3%
Ka'upulehu	31.2%	59.9%	5.8%	0.2%	0.0%	1.8%	0.0%	1.3%
Keawaiki	16.7%	74.9%	5.7%	0.4%	0.2%	0.3%	0.1%	1.7%
Kealakekua	28.6%	65.0%	4.1%	0.7%	0.1%	0.1%	0.0%	1.4%
Ke'ei	28.7%	58.4%	3.6%	0.4%	6.9%	0.3%	0.0%	1.7%
Keopuka	15.6%	75.8%	4.1%	1.3%	0.1%	1.8%	0.1%	1.2%
Kualanui	59.8%	33.4%	3.7%	0.1%	1.3%	0.3%	0.1%	1.1%
Lapakahi	11.4%	56.7%	1.6%	0.9%	0.2%	28.6%	0.1%	1.4%
Makalawena	47.6%	47.5%	1.9%	0.1%	0.0%	1.8%	0.2%	0.6%
Manukā	33.2%	52.9%	9.7%	1.2%	0.1%	1.5%	0.0%	1.5%
N. Keauhou	31.1%	61.4%	5.0%	0.4%	0.4%	0.5%	0.1%	1.1%
Omaka'a	27.1%	61.8%	2.5%	0.2%	0.2%	7.7%	0.0%	0.5%
Papawai	38.3%	39.9%	3.1%	0.6%	4.0%	1.9%	11.0%	1.2%
Puakō	47.8%	42.0%	6.7%	0.6%	0.2%	0.3%	0.0%	2.4%
Red Hill	33.2%	59.4%	2.2%	1.8%	0.1%	2.7%	0.0%	0.6%
S. Oneo	61.9%	31.7%	3.7%	0.9%	0.1%	1.2%	0.1%	0.0%
Waiaka'ilio	42.5%	47.7%	5.5%	0.5%	0.1%	1.2%	0.1%	2.2%
Wawaloli	37.5%	55.3%	3.1%	0.1%	0.0%	0.4%	3.1%	0.5%
Wawaloli Beach	42.3%	52.8%	0.3%	0.0%	0.0%	3.1%	0.3%	1.3%
Keauhou X	57.6%	37.6%	3.3%	0.3%	0.1%	0.6%	0.5%	0.3%
Keauhou Y	40.3%	55.0%	3.0%	0.1%	0.2%	1.0%	0.0%	0.4%
Keauhou Z	42.5%	45.9%	6.6%	0.4%	0.1%	1.2%	2.6%	0.7%
Okoe Bay	34.0%	55.3%	6.0%	0.0%	0.1%	3.3%	0.0%	1.3%
Old Kona Airport	53.2%	25.2%	2.4%	0.3%	0.1%	10.8%	8.0%	0.0%
Unualoha	36.8%	57.3%	1.1%	0.1%	1.1%	1.4%	0.3%	1.8%

Appendix E. West Hawai'i Coral Cover By Species 2007 Surveys

Survey Site	<i>Montipora capitata</i>	<i>Montipora patula</i>	<i>Pavona varians</i>	<i>Pocillopora meandrina</i>	<i>Porites compressa</i>	<i>Porites evermanni</i>	<i>Porites lobata</i>	Other
Anaeho'omalu	1.1%	0.8%	0.6%	0.3%	14%	0.0%	14.2%	0.4%
Ho'okena	3.5%	0.2%	0.0%	2.3%	3.4%	0.4%	19.4%	0.0%
Honokōhau	0.2%	0.0%	0.0%	0.3%	16.0%	0.5%	31.4%	0.4%
Kalahiki	0.1%	0.0%	0.0%	0.3%	13.6%	1.0%	25.5%	0.1%
Kamilo	0.6%	0.1%	0.1%	0.0%	17.1%	0.0%	20.5%	0.2%
Ka'upulehu	0.2%	0.1%	0.1%	0.2%	2.8%	0.0%	27.6%	0.2%
Keawaiki	0.7%	1.6%	1.9%	0.0%	7.4%	0.0%	4.9%	0.1%
Kealakekua	0.1%	0.0%	1.1%	0.2%	11.9%	0.2%	14.9%	0.4%
Ke'ei	0.2%	0.0%	0.1%	0.1%	20.2%	1.5%	6.7%	0.0%
Keopuka	0.2%	0.3%	0.0%	4.8%	1.6%	0.6%	8.2%	0.1%
Kualanui	0.1%	0.4%	0.1%	0.7%	3.2%	18.7%	36.8%	0.0%
Lapakahi	0.1%	0.0%	0.0%	0.0%	1.7%	0.2%	9.4%	0.1%
Makalawena	2.0%	2.6%	1.7%	1.5%	6.2%	0.1%	27.8%	5.7%
Manukā	0.3%	0.0%	0.0%	0.3%	9.9%	1.0%	22.2%	0.1%
N. Keauhou	0.0%	0.0%	0.3%	0.0%	21.2%	0.1%	9.7%	0.0%
Keauhou X	0.1%	0.0%	0.1%	0.0%	18.8%	2.9%	35.7%	0.0%
Keauhou Y	0.1%	0.1%	0.0%	0.0%	26.1%	0.0%	13.2%	0.0%
Keauhou Z	0.0%	0.0%	0.0%	0.1%	24.0%	0.5%	19.1%	0.0%
Okoe Bay	0.3%	0.0%	0.0%	0.3%	4.5%	2.6%	26.3%	0.0%
Old Kona Airport	0.2%	0.0%	0.0%	0.5%	14.1%	0.6%	38.0%	0.0%
Omaka'a	0.9%	0.1%	0.1%	1.2%	7.7%	2.3%	14.9%	0.1%
Papawai	0.2%	0.1%	0.1%	0.4%	3.5%	1.8%	32.4%	0.3%
Puakō	1.3%	0.7%	0.2%	0.5%	17.2%	0.3%	27.2%	1.0%
Red Hill	1.0%	0.2%	0.1%	1.4%	10.2%	1.7%	19.4%	0.0%
S. Oneo	0.2%	0.5%	0.6%	0.3%	30.5%	1.7%	28.2%	0.0%
Unualoha	1.0%	0.1%	0.0%	3.3%	4.5%	0.3%	26.5%	0.2%
Waiaka'ilio	0.5%	0.4%	0.5%	0.3%	14.7%	0.0%	26.4%	0.1%
Wawaloli	0.5%	0.0%	0.1%	3.9%	4.0%	0.3%	28.0%	0.9%
Wawaloli Beach	1.1%	0.0%	0.0%	0.9%	4.0%	1.4%	34.8%	0.0%

Appendix F. West Hawai'i Benthic Cover 2003 Surveys

Broad Benthic Categories

Survey Site	Coral	Turf-Bare	Crustose Coralline	NCC Macroalgae	Macroalgae	Sand	Sessile Invert	Other
'Anaeho'omalu	41.2%	38.8%	8.6%	0.6%	0.0%	3.3%	0.0%	7.5%
Ho'okena	28.5%	55.3%	6.1%	4.3%	0.2%	1.0%	0.3%	4.3%
Honokōhau	48.3%	18.5%	6.8%	0.5%	0.1%	1.7%	11.6%	12.4%
Kalahiki	37.1%	45.6%	5.4%	2.8%	0.3%	3.1%	0.0%	5.7%
Kamilo	49.5%	29.1%	7.4%	3.9%	1.2%	1.1%	0.0%	7.9%
Ka'upulehu	40.9%	40.7%	8.5%	0.3%	0.0%	1.6%	0.0%	7.9%
Keawaiiki	29.9%	51.7%	9.4%	0.0%	0.6%	0.2%	0.0%	8.1%
Kealakekua	27.7%	51.1%	8.0%	2.5%	0.0%	0.0%	0.0%	10.7%
Ke'ei	31.3%	40.0%	14.3%	0.9%	0.0%	0.2%	0.0%	13.4%
Keopuka	16.5%	62.5%	8.2%	1.8%	0.0%	1.3%	0.0%	9.6%
Kualanui	53.3%	36.0%	4.6%	0.7%	0.0%	0.4%	0.2%	4.7%
Lapakahi	19.5%	53.8%	1.4%	0.9%	0.0%	23.0%	0.0%	1.3%
Makalawena	45.2%	44.8%	4.0%	0.3%	0.0%	2.3%	0.1%	3.3%
Manukā	30.8%	50.4%	9.0%	2.7%	0.1%	2.1%	0.0%	4.8%
N. Keauhou	32.9%	41.5%	15.1%	0.4%	0.0%	0.2%	1.3%	8.5%
Omaka'a	30.2%	52.2%	4.2%	0.7%	0.0%	8.4%	0.0%	4.3%
Papawai	32.8%	30.1%	6.2%	0.5%	0.0%	3.0%	19.8%	7.6%
Puakō	49.9%	32.2%	7.5%	0.9%	0.0%	0.9%	0.0%	8.6%
Red Hill	31.5%	40.9%	6.6%	3.9%	0.2%	5.3%	0.8%	10.7%
S. Oneo	57.0%	23.3%	10.5%	0.3%	0.1%	2.1%	0.2%	6.6%
Waiaka'ilio	54.4%	29.1%	5.3%	0.9%	0.8%	1.3%	0.1%	8.1%
Wawaloli	37.9%	45.8%	2.3%	0.2%	0.3%	2.0%	2.5%	9.0%
Wawaloli Beach	33.8%	51.9%	2.4%	0.2%	0.0%	7.1%	0.3%	4.3%

Appendix G. West Hawai'i Coral Cover By Species 2003 Surveys

Survey Site	<i>Montipora capitata</i>	<i>Montipora patula</i>	<i>Pavona varians</i>	<i>Pocillopora meandrina</i>	<i>Porites compressa</i>	<i>Porites evermanni</i>	<i>Porites lobata</i>	Other
'Anaeho'omalu	0.8%	2.2%	1.0%	1.1%	15.2%	0.2%	19.6%	1.2%
Ho'okena	1.6%	0.7%	0.0%	2.0%	2.0%	0.3%	19.3%	2.4%
Honokōhau	0.0%	0.0%	0.0%	0.2%	14.4%	1.8%	31.8%	0.0%
Kalahiki	0.0%	0.0%	0.0%	0.2%	13.4%	0.0%	22.9%	0.6%
Kamilo	0.8%	0.2%	0.1%	0.2%	23.3%	0.1%	24.3%	0.4%
Ka'upulehu	0.2%	0.1%	0.0%	0.3%	6.7%	1.1%	31.9%	0.4%
Keawaiki	0.5%	3.8%	1.4%	0.9%	12.7%	0.0%	8.9%	1.6%
Kealakekua	0.1%	0.3%	1.9%	0.2%	10.6%	0.0%	13.7%	0.8%
Ke'ei	0.1%	0.0%	0.1%	0.0%	19.6%	1.8%	9.4%	0.1%
Keopuka	0.0%	0.1%	0.1%	4.2%	1.0%	0.0%	9.6%	1.6%
Kualanui	0.5%	0.5%	0.1%	0.1%	3.0%	13.7%	34.3%	1.2%
Lapakahi	0.2%	0.0%	0.0%	0.6%	3.1%	0.0%	15.4%	0.1%
Makalawena	1.0%	4.0%	1.0%	1.0%	6.4%	0.5%	26.5%	4.7%
Manukā	0.2%	0.0%	0.0%	0.4%	7.6%	0.4%	21.5%	0.7%
N. Keauhou	0.0%	0.0%	0.6%	0.0%	16.2%	0.0%	15.0%	1.0%
Omaka'a	0.5%	0.4%	0.1%	0.2%	6.8%	2.3%	18.4%	1.4%
Papawai	0.2%	0.1%	0.0%	0.8%	1.8%	0.8%	28.1%	1.0%
Puakō	0.4%	1.7%	0.3%	0.7%	16.9%	0.2%	28.5%	1.3%
Red Hill	0.6%	0.1%	0.1%	0.6%	10.0%	2.0%	16.9%	1.1%
S. Oneo	0.2%	0.6%	0.4%	0.2%	27.2%	1.9%	25.4%	1.0%
Waiaka'ilio	0.6%	2.3%	0.1%	0.7%	19.4%	0.0%	30.5%	0.8%
Wawaloli	0.1%	0.1%	0.0%	5.5%	3.5%	0.0%	27.3%	1.3%
Wawaloli Beach	0.4%	0.1%	0.0%	1.5%	3.2%	1.7%	26.1%	0.7%

Appendix H. 2003, 2007 and 2011 Octocoral Percent Cover Comparison

Site (North to South)	Year			2003/2007	2003/2011	2007/2011
	2003	2007	2011	P=	P=	P=
Lapakahi (01)	0.0%	0.0%	0.0%	N/A	N/A	N/A
Kamilo (2)	0.0%	0.0%	0.0%	N/A	N/A	N/A
Waiakailio Bay (03)	0.0%	0.0%	0.0%	N/A	N/A	N/A
Puako (4)	0.0%	0.0%	0.0%	N/A	N/A	N/A
Anaehoomalu (05)	0.0%	0.0%	0.0%	N/A	N/A	N/A
Keawaiiki (06)	0.0%	0.0%	0.0%	N/A	N/A	N/A
Kaupulehu (07)	0.0%	0.0%	0.0%	N/A	N/A	N/A
Makalawena (8)	0.0%	0.0%	0.0%	N/A	N/A	N/A
Wawaloli Beach (09)	0.4%	0.3%	1.0%	0.908	0.276	0.212
Wawaloli (10)	2.3%	3.1%	3.6%	0.232	0.324	0.696
Honokohau (11)	10.6%	12.7%	13.0%	0.592	0.838	0.971
Papawai (13)	18.2%	10.9%	6.9%	0.018	0.029	0.137
S. Oneo Bay (14)	0.2%	0.1%	0.0%	0.058	0.092	0.391
N. Keauhou (15)	1.2%	0.1%	0.0%	0.13	0.124	0.391
Kualanui Pt. (16)	0.1%	0.1%	0.0%	0.231	0.058	0.391
Red Hill (17)	0.5%	0.0%	0.0%	0.262	0.300	0.391
Keopuka (18)	0.0%	0.0%	0.0%	N/A	N/A	N/A
Kealakekua Bay (19)	0.0%	0.0%	0.0%	N/A	N/A	N/A
Ke'ei (20)	0.0%	0.0%	0.0%	N/A	N/A	N/A
Hookena (Kalahiki) (21)	0.2%	0.0%	0.0%	0.141	N/A	N/A
Hookena (Auau) (22)	0.0%	0.0%	0.0%	N/A	N/A	N/A
Milolii (Omakaa) (23)	0.0%	0.0%	0.0%	N/A	N/A	N/A
Milolii (Manuka) (24)	0.0%	0.0%	0.0%	N/A	N/A	N/A

APPENDIX I. Individuals per 100 m² by species at East Hawai'i sites (n = 224 transects at Waioape; n = 172 at Richardson's Ocean Center).

Taxa	OPEN	MLCD	ROC
<i>Abudefduf abdominalis</i>	0.09	0.74	3.20
<i>Abudefduf sordidus</i>	0.00	0.22	0.03
<i>Abudefduf vaigiensis</i>	0.00	0.04	0.05
<i>Acanthurus achilles</i>	0.00	0.04	0.01
<i>Acanthurus blochii</i>	0.02	0.00	0.00
<i>Acanthurus leucopareius</i>	0.03	0.25	0.25
<i>Acanthurus nigrofuscus</i>	10.35	2.92	6.18
<i>Acanthurus nigroris</i>	0.00	0.04	0.02
<i>Acanthurus triostegus</i>	1.83	2.48	3.36
<i>Aetobatis narinari</i>	0.00	0.00	0.00
<i>Aluterus scriptus</i>	0.00	0.00	0.00
<i>Anampses chrysocephalus</i>	0.00	0.00	0.00
<i>Anampses cuvier</i>	0.03	0.00	0.01
<i>Apogon kallopterus</i>	0.01	0.00	0.00
<i>Apogon menesemus</i>	0.00	0.00	0.00
<i>Arothron hispidus</i>	0.03	0.02	0.01
<i>Arothron meleagris</i>	0.08	0.24	0.07
<i>Asterropteryx semipunctatus</i>	0.03	0.00	0.00
<i>Aulostomus chinensis</i>	0.05	0.02	0.01
<i>Belonidae</i>	0.00	0.01	0.01
<i>Blenniella gibbifrons</i>	0.02	0.01	0.03
<i>Blenniidae</i>	0.00	0.00	0.00
<i>Bodianus bilunulatus</i>	0.03	0.00	0.00
<i>Calotomus carolinus</i>	0.01	0.00	0.00
<i>Cantherhines dumerilii</i>	0.01	0.01	0.01
<i>Canthigaster amboinensis</i>	0.36	1.18	1.06
<i>Canthigaster jactator</i>	3.65	2.67	1.58
<i>Canthigasteridae</i>	0.00	0.00	0.00
<i>Caracanthus typicus</i>	0.00	0.01	0.04
<i>Centropyge potteri</i>	0.00	0.00	0.00
<i>Cephalopholis argus</i>	0.15	0.41	0.01
<i>Chaetodon auriga</i>	0.10	0.05	0.04
<i>Chaetodon lineolatus</i>	0.00	0.00	0.00
<i>Chaetodon lunula</i>	1.50	2.15	0.70
<i>Chaetodon lunulatus</i>	0.00	0.00	0.04
<i>Chaetodon miliaris</i>	0.03	0.00	0.00
<i>Chaetodon multicinctus</i>	0.00	0.00	0.00
<i>Chaetodon ornatissimus</i>	0.12	0.22	0.03
<i>Chaetodon quadrimaculatus</i>	0.49	0.48	0.44
<i>Chaetodon unimaculatus</i>	0.08	0.00	0.00
<i>Cheilio inermis</i>	0.00	0.00	0.00
<i>Chlorurus perspicillatus</i>	0.38	0.04	0.00
<i>Chlorurus sordidus</i>	16.30	1.93	0.63

<i>Chromis agilis</i>	0.09	0.01	0.14
<i>Chromis ovalis</i>	0.78	0.02	0.00
<i>Chromis hanui</i>	0.00	0.00	0.02
<i>Chromis vanderbilti</i>	10.01	9.33	2.82
<i>Cirrhitops fasciatus</i>	0.01	0.04	0.64
<i>Cirrhitus pinnulatus</i>	0.01	0.13	0.14
<i>Cirripectes vanderbilti</i>	0.89	0.54	0.18
<i>Coris flavovittata</i>	0.00	0.01	0.01
<i>Coris gaimard</i>	0.33	0.47	0.41
<i>Coris venusta</i>	0.02	0.08	0.32
<i>Ctenochaetus hawaiiensis</i>	0.00	0.00	0.00
<i>Ctenochaetus strigosus</i>	0.41	0.31	0.00
<i>Dascyllus albisella</i>	0.19	0.03	0.01
<i>Dendrochirus barberi</i>	0.00	0.00	0.00
<i>Diodon hystrix</i>	0.00	0.00	0.00
<i>Echidna nebulosa</i>	0.02	0.00	0.00
<i>Exallias brevis</i>	0.01	0.04	0.02
<i>Fistularia commersonii</i>	0.33	0.09	0.06
<i>Forcipiger flavissimus</i>	0.01	0.06	0.00
<i>Forcipiger longirostris</i>	0.00	0.02	0.00
<i>Gnatholepis anjerensis</i>	0.00	0.00	0.00
<i>Gomphosus varius</i>	4.88	5.74	1.23
<i>Gymnomuraena zebra</i>	0.01	0.02	0.03
<i>Gymnothorax eurostus</i>	0.00	0.04	0.01
<i>Gymnothorax flavimarginatus</i>	0.02	0.03	0.02
<i>Gymnothorax melatremus</i>	0.00	0.00	0.00
<i>Gymnothorax meleagris</i>	0.01	0.04	0.05
<i>Gymnothorax sp.</i>	0.02	0.01	0.02
<i>Gymnothorax undulatus</i>	0.01	0.00	0.00
<i>Halichoeres ornatissimus</i>	0.04	1.04	0.58
<i>Hemitaurichthys thompsoni</i>	0.00	0.00	0.00
<i>Kyphosus bigibbus</i>	0.00	0.28	0.00
<i>Kyphosus sp.</i>	0.00	0.26	0.00
<i>Kyphosus vaigiensis</i>	0.00	0.04	0.00
<i>Labroides phthirophagus</i>	1.50	0.97	0.05
<i>Lutjanus fulvus</i>	0.02	0.00	0.00
<i>Lutjanus kasmira</i>	0.02	0.40	0.00
<i>Lutjanus sp.</i>	0.00	0.00	0.00
<i>Macropharyngodon geoffroy</i>	0.00	0.05	0.06
<i>Melichthys vidua</i>	0.00	0.01	0.00
<i>Mulloidichthys flavolineatus</i>	0.04	0.01	0.03
<i>Mulloidichthys vanicolensis</i>	0.04	0.00	0.00
<i>Myrichthys magnificus</i>	0.00	0.00	0.00
<i>Naso lituratus</i>	0.07	0.05	0.04
<i>Naso unicornis</i>	0.03	0.00	0.01
<i>Neomyxus leuciscus</i>	0.00	0.01	0.11
<i>Neoniphon sammara</i>	0.01	0.00	0.00
<i>Novaculichthys taeniourus</i>	0.01	0.01	0.02

<i>Ostracion meleagris</i>	0.05	0.19	0.05
<i>Oxycheilinus unifasciatus</i>	0.07	0.02	0.02
<i>Paracirrhites arcatus</i>	0.08	0.19	0.58
<i>Paracirrhites forsteri</i>	0.00	0.04	0.05
<i>Parupeneus bifasciatus</i>	0.15	0.07	0.00
<i>Parupeneus cyclostomus</i>	0.05	0.01	0.01
<i>Parupeneus multifasciatus</i>	0.34	0.02	0.02
<i>Parupeneus porphyreus</i>	0.00	0.03	0.00
<i>Pervagor aspricaudus</i>	0.01	0.00	0.00
<i>Plagiotremus ewaensis</i>	0.04	0.05	0.05
<i>Plagiotremus goslinei</i>	0.34	0.41	0.07
<i>Platybelone argalus</i>	0.00	0.09	0.08
<i>Plectroglyphidodon imparipennis</i>	1.58	2.98	8.22
<i>Plectroglyphidodon johnstonianus</i>	0.96	1.07	1.42
<i>Plectroglyphidodon sindonis</i>	0.00	0.00	0.02
<i>Priolepis aureoviridis</i>	0.00	0.00	0.00
<i>Pseudocheilinus evanidus</i>	0.03	0.02	0.02
<i>Pseudocheilinus octotaenia</i>	0.14	0.06	0.00
<i>Pseudocheilinus tetrataenia</i>	0.13	0.14	0.01
<i>Pseudojuloides cerasinus</i>	0.00	0.00	0.00
<i>Pterois sphex</i>	0.00	0.00	0.00
<i>Rhinecanthus rectangulus</i>	0.04	0.04	0.13
<i>Sargocentron diadema</i>	0.01	0.00	0.00
<i>Sargocentron punctatissimum</i>	0.02	0.00	0.01
<i>Sargocentron xantherythrum</i>	0.00	0.03	0.00
<i>Scarus dubius</i>	0.29	0.02	0.01
<i>Scarus psittacus</i>	11.69	1.85	0.47
<i>Scarus rubroviolaceus</i>	0.62	0.28	0.20
<i>Scuticaria tigrinus</i>	0.00	0.01	0.00
<i>Sebastapistes coniota</i>	0.00	0.06	0.22
<i>Stegastes fasciolatus</i>	3.21	19.10	17.38
<i>Stethojulis balteata</i>	5.04	7.98	14.18
<i>Synodus binotatus</i>	0.00	0.01	0.03
<i>Synodus sp.</i>	0.02	0.02	0.00
<i>Synodus ulae</i>	0.00	0.01	0.00
<i>Synodus variegatus</i>	0.01	0.00	0.00
<i>Taenianotus triacanthus</i>	0.00	0.01	0.00
<i>Thalassoma ballieui</i>	0.02	0.04	0.05
<i>Thalassoma duperrey</i>	36.21	35.76	22.38
<i>Thalassoma purpureum</i>	0.00	0.01	0.01
<i>Thalassoma quinquevittatum</i>	0.02	0.00	0.00
<i>Thalassoma trilobatum</i>	0.04	0.06	0.15
Unidentified 1	0.05	0.08	0.05
<i>Zanclus cornutus</i>	0.06	0.10	0.03
<i>Zebrasoma flavescens</i>	0.35	0.03	0.01
Pooled Individuals/100 m2 =	117.3	107.2	90.7