

GENERAL CLIMATIC CONDITIONS.

By ALFRED J. HENRY, Chief of Division of Meteorological Records.

ATMOSPHERIC PRESSURE.

The numerical values of annual mean pressure for 1899 are given in Tables I and II. The method of reduction to sea level in use during the year was the same as in former years, with the exception that an appropriate correction for variations in the force of gravity with latitude has been applied since January 1, 1899. In other respects the annual mean values are comparable with those of the preceding and other years in which Professor Hazen's method of reduction was used.

In addition to the table of reduced pressures, referred to in the preceding paragraph, a second table has been formed (Table III), in conformity with the custom of previous years, by reducing the actual pressures to sea level and standard gravity in accordance with the tables and methods of the International Meteorological Committee, as explained in the MONTHLY WEATHER REVIEW for 1895, Volume XXIII, pages 492-494. The reduced pressures so obtained appear in Table III and on Chart I. The data in the last column of Table III are the pressures at 10,000 feet above sea level, obtained by assuming a uniform decrement of temperature at the rate of 2° F. per 1,000 feet (0.37° C. per 100 meters), as in former annual summaries; the resulting isobars are shown on Chart II.

The distribution of mean pressure at sea level for 1899 is shown by the isobars on Chart I. In general, the pressure distribution for the year 1899 differs but slightly from that of 1898. Pressure was generally above the normal east of the Mississippi River in both years. It was markedly above the normal over Nova Scotia and the Maritime Provinces of Canada in 1898 and also in 1899, although in a less degree. In the latter year the Atlantic high, as traced by the isobar of 30.05 mean annual pressure, extended several hundred miles farther to the northwestward than was the case in 1898. Pressure on the Pacific coast and Plateau region was slightly lower in 1899 than in 1898.

On the immediate Gulf coast of Louisiana, Mississippi and Alabama pressure was from .02 to .04 inch above normal in both years, while less than 200 miles inland, viz, at Vicksburg, Meridian, and Montgomery pressure was from .01 to .04 inch below normal. In both years pressure was also below normal

from the Texas coast westward to Arizona and southern California. The rainfall of both years was likewise less than the normal amount. In mentioning these facts the writer does not intend to convey the impression that they stand in the relation of cause and effect. The fact that there was an average difference of 0.07 inch in pressure between Vicksburg and New Orleans, 0.04 between Mobile and Montgomery, and the same amount between Atlanta and Jacksonville would seem to suggest rather marked changes in the normal air motions along the Gulf coast.

In the Annual Summary for 1898 attention was called to a trough of low pressure which apparently paralleled the foothills of the Rocky Mountains in that year. A similar trough appears on the pressure chart for the current year and the precipitation generally throughout the axis of the trough was above normal as in 1898.

TEMPERATURE.

Although the year was characterized by some of the coldest weather experienced within the last twenty or thirty years, the average temperature on the whole was above normal.

During the greater part of January there were no severe cold waves, but, beginning with the first week in February, the most remarkable cold wave, or series of cold waves, in the history of the Weather Bureau traversed the United States from the north Pacific to the south Atlantic coasts, damaging crops and fruits in the Southern States to a very great extent. The lowest temperatures on record since the beginning of observations were recorded at a number of points in the North Pacific coast States during the first eight days of the month. From the 9th to the 12th of the month the coldest weather on record was reported at a number of points in the Central, Western, and Northwestern States. During the 13th and 14th a cold wave overspread the Southern and Eastern States attended on the 18th by the lowest temperatures ever recorded at many points in the Southern and Gulf States. March was a cold, wintry month, and the spring was generally backward with much snow and unseasonable weather east of the Rocky Mountains.

TABLE A.—Average monthly and annual departures of temperature from the normal during 1899.

Districts.	Number of stations.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Annual.
New England.....	10	+0.5	-2.7	0.0	+0.9	+0.2	+2.1	+0.3	-0.2	-0.8	+1.9	0.0	+2.6	+0.4
Middle Atlantic.....	12	0.0	-5.6	+0.9	+1.0	+0.9	+1.6	+0.0	+1.1	-0.9	-2.5	+1.6	+0.3	+0.2
South Atlantic.....	10	+0.2	-5.5	+2.3	+2.4	+2.1	+1.5	-0.3	+2.1	0.0	-1.9	+2.2	+1.5	+0.2
Florida Peninsula.....	2	+2.0	-1.2	+1.4	-2.2	+1.4	-0.1	-0.9	+0.7	+0.2	-0.7	+0.2	+0.2	+0.2
East Gulf.....	6	-0.8	-9.3	+1.5	-2.4	+1.3	+1.1	+0.1	+2.0	-0.7	-2.5	+2.0	-1.5	0.0
West Gulf.....	7	+0.6	-10.4	+1.6	-2.0	+2.9	+0.3	0.0	+3.5	+0.3	-3.8	+2.2	-2.1	+0.2
Ohio Valley and Tennessee.....	11	+0.4	-10.6	+0.2	+1.3	+2.4	+1.5	+0.5	+3.2	+0.1	+4.9	+2.5	+2.9	+0.5
Lower Lakes.....	9	+0.2	-5.3	-0.7	+4.3	+1.7	+1.2	-0.3	+2.2	+0.2	+4.9	+2.0	+0.2	+0.5
Upper Lakes.....	10	-0.5	-6.3	-5.0	+3.7	+2.1	+0.9	-0.6	+2.5	+0.3	+5.0	+2.0	+0.1	+0.6
North Dakota.....	3	+4.3	-5.9	-13.6	-2.0	-1.6	-1.1	0.0	-0.2	+0.7	+0.1	+12.5	+0.3	-0.6
Upper Mississippi.....	11	+2.9	-9.7	-6.4	+1.2	+1.6	+1.1	-0.1	+3.1	-0.8	+5.1	+9.1	-1.7	+0.5
Missouri Valley.....	10	+5.0	-9.5	-8.2	+1.6	+2.1	+0.6	-0.5	+3.5	-0.9	+5.2	+8.4	-3.1	+0.2
Northern Slope.....	7	+4.2	-12.2	-10.6	-3.1	+2.6	+1.1	-0.5	+1.3	+0.2	-1.3	+6.5	-0.2	-1.4
Middle Slope.....	3	-3.0	-12.1	+4.3	-0.2	+2.6	+0.4	-1.4	+4.0	+1.2	+3.6	+6.6	-2.6	+0.1
Southern Slope.....	2	+0.8	-3.9	+2.0	-1.0	+1.6	-1.9	-1.8	+6.4	+1.7	+3.2	+4.2	-1.6	+0.4
Southern Plateau.....	5	+0.5	-1.1	+0.1	+1.5	-3.1	-0.8	-0.2	-0.9	+2.9	-1.1	+2.5	+1.1	+0.2
Middle Plateau.....	3	+5.6	-0.6	+1.9	-0.2	-6.4	-0.4	+0.6	-5.6	+2.5	-5.5	+3.9	-2.9	-0.7
Northern Plateau.....	3	+6.3	-4.0	+1.3	-3.0	-5.5	-1.2	+1.2	-6.2	+4.5	-2.0	+7.9	-0.2	-0.3
North Pacific.....	3	+1.9	-1.6	-2.6	-1.6	-4.7	-2.6	+0.1	-2.9	+2.4	-0.8	+5.3	+1.5	-0.5
Middle Pacific.....	5	+2.8	0.0	-1.1	+0.5	-3.5	+0.1	-1.0	-2.6	+1.2	-1.3	+1.9	-0.9	-0.3
South Pacific.....	4	+3.3	-0.2	-1.2	+0.2	-3.8	+0.2	-0.3	-3.1	+1.3	-1.9	+1.1	+0.9	-0.8