

funnel gage (of copper) and a Snowdon funnel gage (of japanned iron) gave almost identical totals, while during the winter the former gave 0.136 inch more per month. The annual total was about 30 inches. During March and April the shallow-funnel gage gave on the average 0.027 inch more per month, during the five summer months 0.035 inch less, and during the remaining winter months as much as 0.150 inch less. Apart from the winter months, when the differences were due mainly to snow, the differences were very small. The relative increase in the catch of the Snowdon gage from the spring to the summer months is associated with the general increase in the intensity of the rainfall. The Snowdon gage had therefore marked advantages in winter and in summer. There was, however, a small defect due to the larger surface of the funnel which resulted in greater loss by evaporation. This was apparent only in March and April, when showers alternating with sunshine are a feature of the weather of the British Isles. Symons was satisfied that the advantages of the Snowdon gage far outweighed this small defect,<sup>3</sup> and this conclusion has been borne out by subsequent experience.

A series of comparative readings were subsequently made by Colonel Ward at Rossiniere, Switzerland, with two copper gages with the rims 1 foot, 6 inches above the ground and 9 inches apart. On 30 days during the period October, 1873, to February, 1875, the precipitation took the form of snow, which lay on the ground as soon as it fell. In each case the depth of snow was less than would fill the deep funnel. On these occasions independent estimates of the precipitation were made by inverting the funnel over undrifted snow and measuring in the ordinary way after melting. The comparative readings for these days were: Shallow-funnel gage, 2.67 inches; Snowdon funnel, 5.56 inches; and the independent methods, 5.43 inches.<sup>4</sup> It was concluded that good estimates of snowfall could be made by the use of the Snowdon funnel gage, and that serious losses resulted with the shallow-funnel gage.

Among the numerous experiments with different types of gages carried out in the British Isles, no other comparative readings appear to have been made with funnels of depths other than those already quoted.<sup>5</sup> Reference should, however, be made to the experiments recently carried out by Mr. A. J. Bamford in Ceylon, using deep-funnel and shallow-funnel gages.<sup>6</sup> These experiments were made with special reference to losses by evaporation and to the use of bottles in gages, and they confirm in general the results already given.

In the experiments made in the British Isles using funnels of different depths, the differences in the catch have not been correlated directly with the strength of the wind, as in America.<sup>7</sup> As a matter of fact some of the largest differences in the catch occurred with high wind, but in general there was little correlation between the differences and the strength of the winds. Possibly this is due to the gages in the British Isles generally being in a reasonably sheltered site, since the standard height of the rim of the gage is 1 foot above the ground.

<sup>3</sup> British Rainfall, 1874, pp. 39-40.

<sup>4</sup> See British Rainfall, 1874, face p. 29.

<sup>5</sup> A summary of the experiments made in the British Isles is given in the article on "The Development of Rainfall Measurement in the Last Forty Years," by Dr. H. R. Mill, British Rainfall, 1900, pp. 23-41, although curiously the question of the depth of the vertical funnel is not mentioned.

<sup>6</sup> The Meteorological Magazine, 1930, pp. 81-87.

<sup>7</sup> MONTHLY WEATHER REVIEW, July 1930, vol. 58, pp. 282-283.

## EARLY OPENING OF THE NEW YORK STATE BARGE CANAL

By J. H. SPENCER

[Weather Bureau Office, Buffalo, N. Y.]

The New York State Barge Canal was officially opened this year on April 6, reported to be the earliest in 103 years. The steamer *William Hengerer* and three barges left Buffalo for New York with bonded wheat on the 7th.

There was practically no ice in Lake Erie after March 29. Navigation opened at Buffalo on April 3, with the arrival of the freighter *Coralia* from Detroit loaded with automobiles. The opening of navigation this year was 12 days earlier than the average.

These events reflect the mildness of the winter in this section of the country.

## "MICHAEL SARS" NORTH ATLANTIC DEEP-SEA EXPEDITION, 1910

Reviewed by KATHERINE B. CLARKE

(Report on the scientific results of the *Michael Sars* North Atlantic deep-sea expedition, 1910. Edited by Sir John Murray and J. Hjort. Vol. 1, Deposit Samples by J. Chumley, pp. 1-12; Physical Oceanography and Meteorology, by B. Helland-Hansen, text pp. 1-115, tables and plates pp. 1-102. Published by the Bergen Museum, Bergen, Norway, 1931.)

This volume is the first in a series on the scientific results of the *Michael Sars* expedition, a series which promises to be an exceedingly valuable contribution to the development of oceanography and associated sciences. Planned chiefly as a biological survey of the North Atlantic, a gratifying amount of geophysical and meteorological data of real value was also obtained. Few persons are better qualified by knowledge and experience to write the discussion of these data than the eminent Norwegian oceanographer, Helland-Hansen.

The section on physical oceanography and meteorology has two major divisions, the text and the tables and plates. The text is further outlined in 10 chapters which cover the following topics: Introduction, sea surface and air, subsurface temperatures, salinities and densities (methods), local variations in general, short-period oscillations, the temperatures in the sea, salinities in the North Atlantic, stability, dynamics of the sea, and current measurements. The text is followed by an ample bibliography.

The chief interest from a meteorological viewpoint lies in the discussions of sea and air temperatures, the interaction of ocean and atmosphere, and diurnal, seasonal, and annual variations. Observations were made from June 3 to August 15, 1910. For purposes of statistical compilation these are divided into four series. On Deutsche Seewarte synoptic charts for each day of the cruise the position of the *Michael Sars* is shown. Accompanying graphs give for each day the meteorological conditions observed on the ship.

A very conspicuous positive correlation between surface temperatures and surface salinity was found. Regional variations were the chief cause for great variations in mean surface temperature. The daily period of surface temperature was not as evident as might be expected, but correlated closely with the amount of cloudiness, being more prominent with slight than with extensive cloudiness. A conclusion previously expressed by Helland-Hansen and Nansen, that variations in surface temperature are primarily the result of displacement of the surface

layers and that the warming and cooling effect of the air upon the sea is of secondary importance, has been substantiated by the results of this expedition. Mean air temperature was found to have a strong tendency to follow mean surface temperature. Only rarely was the difference between the two as much as 2° C. These differences though slight are of the utmost importance in atmospheric processes and necessitate the most careful observations of sea and air temperatures. The air was, as a rule, warmer than the sea by day and cooler by night, but 68 per cent of the averages give air temperatures lower than sea temperatures. Variations in humidity correspond closely to the variations in temperature differences between water and air. Moreover, variations in relative humidity coincide almost exactly with variations in absolute humidity.

In the study of gain and loss of heat the author reaches the same conclusions as Harvey;<sup>1</sup> namely, that evaporation is the chief cause for the loss of heat from the sea, excess of outward radiation, and direct convection to the air being of secondary importance; and that the stronger the heating of the surface layers the less the heating of the

subsurface layers, due to greater stability and hence lessened convection to lower levels.

The study of seasonal variations was hampered by the fact that the author could find in literature only three cases where serial observations for deep offshore parts of the eastern North Atlantic have been taken at different seasons at the same geographical position. The author finds on the whole fairly good indication of "direct agreement between annual variations of the surface temperature toward the end of the winter and the temperatures for a considerable time (several months) afterward at 50 and 100 meters," p. 61.

Throughout the volume Helland-Hansen has amplified his discussion with results of his other observations, particularly those made in the North Atlantic in collaboration with F. Nansen. It is regrettable that the publication of results of such expeditions are so delayed. Such delays may only be compensated for by thorough and careful studies of results, such as are exhibited in this publication. The whole volume serves to emphasize the necessity for cooperation between investigators in physical oceanography and meteorology and for the collection of more detailed and accurate marine meteorological data.

<sup>1</sup> Harvey, M. A. Evaporation and Temperature Changes in the English Channel. *J. Marine Biol. Assoc.*, v. 13, pp. 67.8-69.2, 1925

## BIBLIOGRAPHY

C. FITZHUGH TALMAN, in charge of Library

### RECENT ADDITIONS

The following have been selected from among the titles of books recently received as representing those most likely to be useful to Weather Bureau officials in their meteorological work and studies:

**Ångström, Anders.**

Das Ångström-Pyranometer, Type 1930. p. 526-534. figs. 24 cm. (Strahlentherapie, 39. Bd. (1931).)

**Bider, Max.**

Ergebnisse der Beobachtungen mit dem Davoser Frigorimeter in Basel. p. 541-564. figs. 24 cm. (Strahlentherapie, 39. Bd. (1931).)

**Bouasse, H., & others.**

Tourbillons, forces acoustiques, circulations diverses. Tome 1. Paris. 1931. xxiv, 422 p. figs. 25½ cm.

**Davos. Physikalisch-meteorologisches Observatorium.**

Neukonstruktion des "Davoser Frigorimeters." [Davos.] 1931. 2 p. illus. 34½ cm. [Manifolded.]

**Field, R. H.**

Aneroid barometer and altimeter, their characteristics and use in mapping. With an appendix: The field use of the aneroid barometer, by G. C. Cooper. Ottawa. 1931. 36 p. illus. 25 cm. (Canada, dept. inter. Topographic survey. Bul. no. 63.)

**Gincstous, G.**

Récupération de l'humidité atmosphérique. Alger. 1929. 7 p. 24½ cm. (Extr.: Rev. agric. de l'Afrique du Nord.)

**Kirby, Harold Lewis.**

Analysis of meteorology as related to the operation of aircraft. Los Angeles. [c1930.] 120, 14 p. front. illus. diags. 26½ cm.

**McAdie, Alexander.**

Clouds and the airman. p. 49-59. illus. 30 cm. (Nat. aeron. mag. Dec. 1930.)

**Sievert, Otto.**

Wetterkunde. Eine Anleitung zu Wetterverständnis und Wettervoraussage. Dritte vermehrte Aufl. Berlin. [1930.] 168 p. figs. plate (fold.). 21½ cm.

**Talman, Charles Fitzhugh.**

Realm of the air; a book about weather. Indianapolis. [c1931.] x, 318 p. front. plates. 23 cm.

**Vallaux, Camille.**

Les dérives antarctiques en surface et en profondeur. Monaco. n. d. 16 p. figs. 25½ cm. (Bull. de l'Inst. océan., No. 567. 10. fév. 1931.)

**Van Orman, W. T.**

Preliminary meteorological survey for airship bases on the middle Atlantic seaboard. n. p. 1931. v. p. plates. 28½ cm. (For International Zeppelin transport corporation, 52 Wall St., New York City.) [Typewritten.]

**Wagner, A.**

Klimatologie der freien Atmosphäre. Berlin. 1931. 70 p. figs. 27 cm. (Handbuch der Klimatologie. Bd. I, Teil F.)

**Waitz, Karl.**

Die kosmischen Ursachen des Wetters. Leipzig. [c1930.] vii, 95 p. figs. 23½ cm.